



ADAMA

# MAXENTIS®

Provides protection against the most damaging diseases in peas, lentils and canola. It features two distinct, elite modes of action for resistance management (Groups 3 & 11) along with long-lasting systemic activity.



FUNGICIDES

POWERED BY

**Asorbital® 2**

FORMULATION TECHNOLOGY

**GROUPS 3 & 11**

## Active Ingredients

Azoxystrobin 120 g/L + Prothioconazole 90 g/L as an EC

## Packaging

Case: 2 x 8.45 L jugs  
Drum: 118.1 L

## Application Rates and Acres Treated

### CANOLA

Rate: 443 ml/ac

Acres Treated:

19 ac/jug

267 ac/drum

### PEAS, LENTILS & SOYBEANS

Rate: 422ml/ac

Acres Treated:

20 ac/jug

280 ac/drum

## KEY BENEFITS

- Unique combination of prothioconazole and azoxystrobin
- Enhanced EC formulation – with bulk packaging capabilities
- Multi-mode activity for resistance management (Groups 3 & 11)
- Broad-spectrum disease control with protective and curative action
- Whole plant protection: translaminar and systemic movement
- Wide window of application

## REGISTERED CROPS\*

- Canola
- Peas
- Lentils
- Soybeans

## KEY DISEASES CONTROLLED (OILSEEDS AND PULSES)\*

- Anthracnose
- Mycosphaerella blight (suppression)
- White mould (*Sclerotinia*)

## HOW IT WORKS

Two powerful active ingredients, combined for the first time in Canada, that provide protective and curative action on a wide range of diseases, across multiple crops.

\* See label for the complete list of registered crops as well diseases controlled and rates for each crop.



# MAXENTIS®

## APPLICATION TIMING AND CROP STAGING (PULSES AND OILSEEDS)

Crop	Diseases	Rate (ml/ac)	Timing
Canola	White mould ( <i>Sclerotinia</i> )	443	Apply when the crop is in the 20–50% bloom stage. Best protection will be achieved when the fungicide is applied prior to petals beginning to fall and will allow for the maximum number of petals to be protected.  Maximum applications per year: 1
Peas	Mycosphaerella blight (suppression) White mould ( <i>Sclerotinia</i> )	422	Apply as a preventive foliar spray at the beginning of flowering or at first sign of disease.  After the initial application, one additional application may be made 10–14 days afterwards if conditions remain favourable for continued or increased disease development. Apply the higher rate when conditions favour disease development, or when growing less disease resistant varieties.  Maximum applications per year: 2
Lentils	Anthraxnose, white mould		
Soybeans	White mould ( <i>Sclerotinia</i> )		

### MIXING INSTRUCTIONS

1. Fill the clean spray tank  $\frac{3}{4}$  full of clean water.
2. Add the required amount of MAXENTIS® and agitate thoroughly.
3. Continue agitation and add the required amount of the tank-mix partner.
4. Complete filling tank to the desired level with water.

### WATER VOLUME

**Ground:** 40 L/ac (10 US gal/ac)

**Aerial:** 20 L/ac (5 US gal/ac)

### RAINFASTNESS

Avoid applying when rain is forecast.

### GRAZING RESTRICTIONS

Forage, hay: 30 days

Grazing or green feed: 6 days

**NOTE:** Do not feed dried pea vines to livestock.

### CROP ROTATIONS

No restrictions

### PRE-HARVEST INTERVALS

- Peas, lentils: 15 days
- Soybeans: 20 days
- Canola: 36 days

**NOTE:** Refer to the label for pre-harvest intervals for all other registered crops.

### STORAGE

Do not freeze.

### WANT TO KNOW MORE?

Click here for full product information including label and SDS documents.



## Quick Tips

Environmental conditions are a major driver of disease pressure and severity. Under high disease pressure conditions a rate of MAXENTIS® at the upper end of the rate range is recommended.

Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. 1.855.264.6262 ADAMA.COM



ADAMA

®/™ MAXENTIS is a registered trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.

© 2024 ADAMA Agricultural Solutions Canada Ltd.