

Active Ingredients:

Fluroxypyr 59.7 g/L Clopyralid 40.00 g/L MCPA 2 EH Ester 240.5 g/L = EC

Application Rates and Acres Treated:

Rate:

Low: 730 ml/ac High: 970 ml/ac

Acres Treated: 11-15 ac/jug

120-160 ac/drum

Packaging:

Case: 2 x 10.93 L jugs Bulk: 116.5 L drums

Water Volume:

Ground: 20 - 40 L/ac

(5-10 US gal/ac)

Aerial: 12-20 L/ac

(3-5 US gal/ac)

Rainfastness:

4 hours

REGISTERED CROPS:

- Spring Wheat
- Durum Wheat
- Winter Wheat

- Spring Barley
- Oats

WEEDS CONTROLLED:

At the low rate of 15 ac/jug will control:

- Annual Sunflowers
- Burdock
- Canada Thistle (low infestations)
- Cleavers
- Cocklebur
- Field Horsetail (top growth)
- Flixweed
- Kochia
- Lamb's Quarters
- Plantain (top growth)
- Prickly Lettuce

- Ragweeds
- Shepherd's Purse
- Stinkweed
- Stork's Bill
- Wild Buckwheat
- Wild Mustard
- Wild Radish
- Vetch
- Volunteer Flax
- Volunteer Sunflowers

At the high rate of 11 ac/jug will control:

- Annual Sow Thistle
- Canada Thistle (medium to high infestations, season long control)
- Chickweed
- · Common Groundsel
- Dandelions (spring rosettes only)
- Hempnettle
- Perennial Sow Thistle (season long control)
- · Redroot Pigweed

- Roundleaf Mallow
- Russian Pigweed
- Scentless Chamomile
- Smartweed
- Tartary Buckwheat
- Volunteer Canola



ESTEEM ALL IN™

HOW IT WORKS:

Moves within the plant to control exposed and underground plant tissues. It mimics naturally occurring plant hormones which control weeds by disrupting normal plant growth patterns. Symptoms of effect include epinasty (twisting of the stems) and swollen nodes.

CROP STAGING:

Spring wheat, durum wheat, spring barley and oats: 3-leaf to just before the flag leaf. Winter wheat in the spring: 3 tiller stage to just before the flag leaf.

REGISTERED AND SUPPORTED TANK MIXES:

- Assert® 300 SC
- BRAZEN ALL IN™
- BISON® 400 L
- BroadBand®
- BRAZEN® II
- LADDER ALL IN®
- Puma® Advance
- Simplicity™ GoDRI™*
- Traxos[®]
- Varro®

*Only use 30 ac/case rate when mixing with Simplicity™ GoDri™

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

- 1. Fill sprayer tank ½ full of water.
- 2. Start sprayer tank agitation.
- 3. Add any dry formulation tank mix partners.
- Add the required amount of grassy weed tank mix partner.
- **5.** Add the required amount of ESTEEM ALL IN^{TM} .
- Add any required adjuvants of surfactants for tank mix partners.
- 7. Complete filling tank with water.

STORAGE:

Do not freeze.

PRE-HARVEST INTERVAL:

Wheat, barley or oats: 60 days

Forage: 7 days

GRAZING RESTRICTIONS:

Do not cut or graze treated fields of wheat, barley or oats within 7 days after application.

Withdraw meat animals from treated fields at least 3 days before slaughter.



Always read and follow label directions.
Toll-free:1.855.264.6262 | Website: ADAMA.COM

BISON®, BRAZEN® II, ESTEEM® and LADDER ALL IN® are registered trademarks, and BRAZEN ALL IN™ is a trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.

© 2024 ADAMA Agricultural Solutions Canada Ltd.

CROP ROTATIONS:

Fields previously treated with ESTEEM ALL IN[™] can be seeded to wheat, barley, oats and rye (not underseeded to forage legumes, clover or alfalfa), canola, field peas, flax, forage grasses, mustard, or can be summer-fallowed

For field peas: a rainfall of 140 mm (5.5 inches) between herbicide application and August 31, and an annual precipitation greater than 175 mm (6.9 inches) is required.



Quick Tips

Flexible rates available - choose what works best for you: 23ac/case for heavy weed pressure OR hard to control weeds; 30ac/case for standard weed pressure.