



DAVAI® Family of Products for Grassy and Broadleaf
Weed Protection

DAVAI® Q PLUS

Excellent rotation partner with dual modes of action for controlling stubborn clumping grasses, such as Persian Darnel and Barnyard Grass, and broadleaf weeds.

Kev Benefit

• DAVAI® Q PLUS combines both Group 1 and Group 2 modes of action for resistance management. DAVAI® Q PLUS has 32% more quizalofop in the case then the competitive product.

Active Ingredient

- Imazamox 80 g/L = SL
- Quizalofop-P-ethyl 100 g/L = EC

Application Rates and Acres Treated:

Rate: 100 mL/ac DAVAI® 80 SL + 194 mL/ac ADAMA Quizalafop + ADAMA MSO @.5% v/v

Acres Treated: 40 ac/case

CLICK HERE FOR FULL **PRODUCT** DETAILS



Best for controlling:

FOXTAIL BARLEY

VOLUNTEER CEREALS (EXPANDED WINDOW)

WINTER ANNUAL GRASSES (DOWNY BROME & JAPANESE BROME)

DAVAI® A PLUS

DAVAI® 80 SL conveniently packaged with ARROW ALL IN® the leading grass control product in pulses.

Key Benefit

• DAVAI® A PLUS combines both Group 1 and Group 2 modes of action for

Active Ingredient

- Imazamox 80 g/L = SL

Application Rates and Acres Treated:

Rate: 150 ml/ac ARROW ALL IN® + 100 ml/ac

Acres Treated: 40 ac/case

CLICK HERE FOR FULL **PRODUCT** DETAILS



Best for controlling:

WILD OATS

Broadleaf Weeds controlled by all:

COTYLEDON TO 4 LEAF • Green Smartweed

- Cleavers (S)
- Cow Cockle
- Flixweed
- Lamb's Quarters
- Redroot Pigweed
- Shepherd's Purse

DAVAI 80[®] SL

Proven broadleaf and grassy weed control in a convenient package that allows for flexible tank-mix options.

Kev Benefit

 Many options for re-cropping when any DAVAI® product is used at labeled rates.

Active Ingredient

Imazamox 80 g/L = SL

Application Rates and Acres Treated:

Rate: 100 mL/ac DAVAI® 80 SL

Acres Treated: 160 ac/case

CLICK HERE FOR FULL PRODUCT **DETAILS**



Best for controlling:

GREEN FOXTAIL

IF OVERALL GRASSY WEED PRESSURE IS LOW

- Stinkweed
- Stork's Bill
- Wild Buckwheat (S)
- Wild Mustard

*

Crop Rotations the following crop year:

Barley

Chickpeas

Flax

SoybeansClearfield Sunflowers

Canary SeedCanola

CornField Peas

LentilsOats

• Wheat (spring, durum)

If less then 150 mm of rainfall between time of application and August 31, it is strongly recommended to refrain from growing canola, durum wheat or canary seed the following year. Contact your ADAMA representative for details.

Registered Crops and Staging		
CROP	STAGE	
Dry beans, Soybeans	Emergence to 3 expanded trifoliate leaves	
Field Peas	1-6 true leaves	
Imidazolinone-tolerant Lentils	1-9 nodes	

	DAVAI® Q PLUS (IMAZAMOX + QUIZALOFOP)	DAVAI® A PLUS (IMAZAMOX + CLETHODIM)	DAVAI® 80 SL (IMAZAMOX)
WEED STAGE:	2 - EARLY TILLERING (EXCEPT WHERE INDICATED)	1-6 TRUE LEAF	1-4 TRUE LEAF
Barnyard Grass	X	X	Х
Crabgrass (smooth, large)		X ⁴	
Downy Brome	X ₃		
Fall Panicum	X	X	
Foxtail Barley	X ²		
Green Foxtail	Х	X	Х
Japanese Brome Grass	X ₃	S	S
Persian Darnel	X ⁵	X	Х
Proso Millet	X	X	
Quack Grass	S ⁴	S	
Volunteer Barley	X	X	Х
Volunteer Canary Seed	X ⁵	X ⁵	Х
Volunteer Canary Grass	X ⁵	Х	
Volunteer Corn	X ⁴	X ⁴	
Volunteer Oats	X ⁶	X	Х
Volunteer Wheat	Х	Х	Х
Wild Oats	X ^{1.6}	Х	Х
Witch Grass	Х	Х	
Yellow Foxtail	X	Х	Х

S - Suppression | ¹ - 1-5 leaf to early tillering; ² - 2-4 leaf + 3 tillers; ³ - 2-5 leaf; ⁴ - 2-6 leaf; ⁵ - 1-4 leaf; ⁶ - Best results on volunteer oats/wild oats if application is made before tillering begins.

