



ADAMA DICAMBA 480 SL

A liquid formulation that offers a convenient tank mix partner for pre-seed weed control ahead of cereals, helping combat Group 2 & 9 resistant weed species.

GROUP 4

Active Ingredient:

Dicamba 480 g/L = SL

Application Rates & Acres Treated:

Rate: 100 ml/ac

Rate when tank mixed with PRIORITY® HL for pre seed/fall application

Acres Treated: 200 ac/case;
100 ac/jug

Packaging:

Case: 2 x 10 L

Water Volume:

Ground: 40 L/ac (10 US gal/ac)

Aerial: Do not apply

Rainfastness:

30 minutes

REGISTERED CROPS:

- Oats
- Spring Barley
- Spring Wheat

WEEDS CONTROLLED:

Weeds controlled when tank mixed with PRIORITY® HL (8 ml/ac) and glyphosate (0.5 REL/ac).

Broadleaf weeds controlled (2-4 leaf stage):

- | | | |
|--------------------------------|---|---------------------------------|
| • Canada Fleabane ² | • Kochia ¹ | • Smartweed |
| • Cleavers | • Lady's Thumb | • Stinkweed |
| • Common Chickweed | • Lamb's Quarters | • Volunteer Canola ⁴ |
| • Common Ragweed ² | • Narrow-Leaved Hawk's Beard ² | • Volunteer Flax |
| • Cow Cockle | • Redroot Pigweed | • Wild Buckwheat ⁵ |
| • Dandelion ³ | • Russian Thistle | • Wild Mustard |
| • Flixweed | • Shepherd's Purse | |
| • Hemp-Nettle | | |

Grassy weeds controlled:

- | | | |
|-----------------|--------------------|-------------------|
| • Downey Brome | • Persian Darnel | • Volunteer Wheat |
| • Giant Foxtail | • Volunteer Barley | • Wild Oats |
| • Green Foxtail | | |

Additional weeds controlled when tank mixed with PRIORITY® HL (8 ml/ac) and glyphosate (1-2.8 REL/ac).

Controlled (2-4 leaf stage):

- | | |
|-------------------------------|--------------------------------------|
| • Annual Sow Thistle | • Quack Grass |
| • Canada Thistle ⁷ | • Perennial Sow Thistle ⁶ |

¹ Control of Group 2, 4 and 9 resistant weeds

² Less than 3 inches in height.

³ Mature plants up to 12 inches in diameter, rosettes, and seedlings.

⁴ Including all herbicide-tolerant canola varieties.

⁵ Up to 5 leaves.

⁶ Applications made at advanced stages will be less effective.

⁷ Rosette stage.

ADAMA DICAMBA 480 SL

HOW IT WORKS:

ADAMA DICAMBA 480 SL controls broadleaf weeds in cereals, field corn, reduced tillage (prior to seeding and reduced tillage fallow), pastures and rangeland grasses, crop-free land (summer fallow and stubble), red fescue, canary seed (*Phalaris canariensis*), seedling grasses grown for seed and forage and low bush blueberries.

APPLICATION TIMING:

Apply prior to seeding wheat, barley and oats. Do not apply later than 2 hours post seeding.

REGISTERED & SUPPORTED TANK MIXES:

- PRIORITY® HL
- Glyphosate

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

1. Fill the clean spray tank ½ full of clean water. Start agitation system.
2. Add the required amount PRIORITY® HL. Continue to agitate.
3. Add the required amount of ADAMA DICAMBA 480 SL. Continue to agitate.
4. Add required amount of glyphosate.
5. Complete filling with remaining amount of water and continue agitation.

GRAZING RESTRICTIONS:

- DO NOT permit lactating dairy animals to graze fields within 7 days after application.
- DO NOT harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions and grazing limitations for each product used in the tank-mix.

CROP ROTATIONS:

When applied prior to August 1st: barley, canola, oats, field peas and wheat (spring, durum, winter) can be seeded the following year.

PRE-HARVEST INTERVALS:

Do not harvest the treated crop within 60 days of application.

STORAGE:

- Store ADAMA DICAMBA 480 SL in its original container only, away from other pesticides, fertilizer, food, or feed.
- Keep the container closed to prevent spills and contamination.
- Keep packages dry at all times.



ADAMA

Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

PRIORITY® is a registered trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

Compatible with all forms of glyphosate.