

ADAMA DICAMBA 480 SL

A liquid formulation that offers a convenient tank mix partner for pre-seed weed control ahead of cereals, helping combat Group 2 & 9 resistant weed species.

Active Ingredient:

Dicamba 480 g/L = SL

Application Rates & Acres Treated:

Rate: 100 ml/ac

Rate when tank mixed with PRIORITY[®] HL for pre seed/fall application

Acres Treated: 200 ac/case; 100 ac/jug

Packaging:

Case: 2 x 10 L

Water Volume:

Ground: 40 L/ac (10 US gal/ac) Aerial: Do not apply

Rainfastness:

30 minutes

GROUP 4

REGISTERED CROPS:

- Oats
- Spring Barley
- Spring Wheat

WEEDS CONTROLLED:

Weeds controlled when tank mixed with PRIORITY® HL (8 ml/ac) and Glyphosate (0.5 REL/ac).

Kochia¹

Broadleaf weeds controlled (2-4 leaf stage):

- Canada Fleabane²
- Cleavers
- Common Chickweed
- Common Ragweed²
- Cow Cockle
- Dandelion³
- Flixweed
- Hemp-Nettle

Grassy weeds controlled:

- Downey Brome
- Giant Foxtail
- Green Foxtail

- Smartweed Lady's Thumb Stinkweed
- Lamb's Quarters Narrow-Leaved
- Hawk's Beard²
- Redroot Pigweed
- Russian Thistle
- Shepherd's Purse
- Persian Darnel
- Volunteer Barley

Volunteer Canola⁴

Wild Buckwheat⁵

• Volunteer Flax

Wild Mustard

- Volunteer Wheat
- Wild Oats

Additional weeds controlled when tank mixed with PRIORITY® HL (8 ml/ac) and glyphosate (1-2.8 REL/ac).

Controlled (2-4 leaf stage):

- Annual Sow Thistle
- Canada Thistle⁷

- Quack Grass
- Perennial Sow Thistle⁶

¹Control of Group 2, 4 and 9 resistant weeds

²Less than 3 inches in height. ³ Mature plants up to 12 inches in diameter, rosettes, and seedlings.

- ⁴Including all herbicide-tolerant canola varieties.
- ⁵Up to 5 leaves.
- ⁶Applications made at advanced stages will be less effective.

⁷Rosette stage.

ADAMA DICAMBA 480 SL

HOW IT WORKS:

ADAMA DICAMBA 480 SL controls broadleaf weeds in cereals, field corn, reduced tillage (prior to seeding and reduced tillage fallow), pastures and rangeland grasses, crop-free land (summer fallow and stubble), red fescue, canary seed (*Phalaris canariensis*), seedling grasses grown for seed and forage and low bush blueberries.

APPLICATION TIMING:

Apply prior to seeding wheat, barley and oats. Do not apply later than 2 hours post seeding.

REGISTERED & SUPPORTED TANK MIXES:

- PRIORITY® HL
- Glyphosate

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

- 1. Fill the clean spray tank ½ full of clean water. Start agitation system.
- 2. Add the required amount PRIORITY® HL. Continue to agitate.
- 3. Add the required amount of ADAMA DICAMBA 480 SL. Continue to agitate.
- 4. Add required amount of Glyphosate.
- 5. Complete filling with remaining amount of water and continue agitation.

GRAZING RESTRICTIONS:

- DO NOT permit lactating dairy animals to graze fields within 7 days after application.
- DO NOT harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions and grazing limitations for each product used in the tank-mix.

STORAGE:

- Store ADAMA DICAMBA 480 SL in its original container only, away from other pesticides, fertilizer, food, or feed.
- Keep the container closed to prevent spills and contamination.
- Keep packages dry at all times.

CROP ROTATIONS:

*For PRIORITY[®] HL, ADAMA DICAMBA 480 SL, & Glyphosate tank mix.

When applied prior to August 1st: barley, canola, oats, field peas and wheat (spring & durum) can be seeded the following year.

When applied after August 1st: only wheat (spring & durum), barley and oats can be seeded the following spring.

PRE-HARVEST INTERVALS:

Do not harvest the treated crop within 60 days of application.



Always read and follow label directions. Toll-free: 1.855.264.6262 | Website: ADAMA.COM

 $\label{eq:product} PRIORITY^{\circ} \ is a registered trademark of an ADAMA Group Company. All other products are $$ trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd. $$$

Compatible with all forms of glyphosate.

QUICK TIPS: