

2025-2245
2025-07-03

(Container)

GROUP 2	HERBICIDE
----------------	------------------

GORDEX™ A

Suspension

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING PEACE RIVER REGION) ONLY

A selective herbicide for postemergent control or suppression of annual broadleaved weeds including cleavers, wild buckwheat, and chickweed in spring wheat (including durum), spring barley, and oats (tank mixes only).

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: florasulam 200 g/L
Contains 1,2-benzisothiazolin-3-one at 0.050% as a preservative

REGISTRATION NO. 35555 PEST CONTROL PRODUCTS ACT

CAUTION



POISON

MAY CAUSE MILD EYE IRRITATION

NET CONTENTS: 0.5 – 1050 L

For emergency medical help and health/safety inquiries call ProPharma Group at 1.877.250.9291 (24 hours a day)

For spill, leak or fire call INFOTRAC at 1.800.535.5053 (24 hours a day)

ADAMA Agricultural Solutions Canada Ltd.
300 – 191 Lombard Avenue
Winnipeg, Manitoba
R3B 0X1
1-855-264-6262

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. DO NOT APPLY BY AIR.

Harmful if inhaled. Avoid inhaling/breathing sprays or mists. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab. **At completion of spraying or end of the day:** Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

CAUTION: Users of **GORDEX™ A** should check weather conditions prior to application of **GORDEX™ A** on crops. Crops can be more sensitive to herbicide application under certain stressful environmental conditions. For example, heavy rain after **GORDEX™ A** application that leaches the active ingredient florasulam into the root zone may cause crop injury, such as yellowing and stunting. In most cases, crops will outgrow the symptoms, but in severe situations reduced crop stand, yield, quality or delayed maturity may occur.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take the container label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

2025-2245
2025-07-03

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

Store this product away from food or feed. Store in original containers in a secure, dry heated storage. **GORDEX™ A** will freeze at -10°C. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Do not contaminate domestic water supplies. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers (containers that can be refilled for the user by the distributor/dealer):

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

Disposal of unwanted, unused product:

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label.

2025-2245
2025-07-03

(Booklet)

GROUP	2	HERBICIDE
--------------	----------	------------------

GORDEX™ A

Suspension

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING PEACE RIVER REGION) ONLY

A selective herbicide for postemergent control or suppression of annual broadleaved weeds including cleavers, wild buckwheat and chickweed in spring wheat (including durum), spring barley, and oats (tank mixes only).

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: florasulam 200 g/L
Contains 1,2-benzisothiazoline-3-one at 0.050% as a preservative

REGISTRATION NO. 35555 PEST CONTROL PRODUCTS ACT

CAUTION



POISON

MAY CAUSE MILD EYE IRRITATION

NET CONTENTS: 0.5 – 1050 L

For emergency medical help and health/safety inquiries call ProPharma Group at 1.877.250.9291 (24 hours a day)

For spill, leak or fire call INFOTRAC at 1.800.535.5053 (24 hours a day)

ADAMA Agricultural Solutions Canada Ltd.
300 – 191 Lombard Avenue
Winnipeg, Manitoba
R3B 0X1
1-855-264-6262

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. DO NOT APPLY BY AIR.

Harmful if inhaled. Avoid inhaling/breathing sprays or mists. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab. **At completion of spraying or end of the day:** Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

CAUTION: Users of **GORDEX™ A** should check weather conditions prior to application of **GORDEX™ A** on crops. Crops can be more sensitive to herbicide application under certain stressful environmental conditions. For example, heavy rain after **GORDEX™ A** application that leaches the active ingredient florasulam into the root zone may cause crop injury, such as yellowing and stunting. In most cases, crops will outgrow the symptoms, but in severe situations reduced crop stand, yield, quality or delayed maturity may occur.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take the container label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

2025-2245
2025-07-03

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

Store this product away from food or feed. Store in original containers in a secure, dry heated storage. **GORDEX™ A** will freeze at -10°C. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Do not contaminate domestic water supplies. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers (containers that can be refilled for the user by the distributor/dealer):

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

Disposal of unused, unwanted product:

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label.

GENERAL INFORMATION

GORDEX™ A is a selective postemergence herbicide for the control or suppression of hard-to-

kill annual broadleaved weeds such as chickweed, wild buckwheat and cleavers in spring wheat (including durum), spring barley, and oats (tank mixes only) not underseeded with legumes. **GORDEX™ A** is mixed with water and applied as a uniform broadcast spray. It is non-corrosive, nonflammable, and nonvolatile.

GORDEX™ A must be applied early postemergence, to the main flush of actively growing broadleaved weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of **GORDEX™ A** by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds. See “DIRECTIONS FOR USE” section of this label for complete use details.

GORDEX™ A stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 2 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

MODE OF ACTION

GORDEX™ A inhibits the production of the ALS enzyme in plants. This enzyme is essential for the production of certain amino acids which are essential for plant growth. **GORDEX™ A** is a Group 2 mode of action herbicide.

GENERAL USE PRECAUTIONS

Sensitive Plants

Do not apply **GORDEX™ A** directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes or tobacco.

Non-Target Sites

Do not apply where proximity of susceptible crops (e.g. canola and legumes) or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Precautions section of the label.

Crop Rotation

Fields previously treated with **GORDEX™ A** can be seeded the following year to barley, canola, peas, wheat or fields can be summerfallowed.

Do not use in successive years at the same site.

This product has potential to leach. Do not apply excessive irrigation.

Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing

2025-2245
2025-07-03

the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label

Spray Equipment Precaution

Do not apply through any type of irrigation system.

To Reduce Spray Drift:

1. Use nozzles delivering higher volumes and coarser droplets.
2. Use low pressures (200 to 275 kPa).
3. Use 100 L/ha of spray solution.
4. Spray when the wind velocity is 15 km/hr or less.
5. Spot treatments should only be applied with a calibrated boom to prevent over-application.

Equipment Clean-Up

Equipment used to apply **GORDEX™ A** should not be used to apply other pesticides to sensitive crops without thorough cleaning. To avoid subsequent injury to crops other than cereals, all spraying equipment must be thoroughly cleaned both inside and out, as follows:

1. Immediately after spraying, drain. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
3. Add 1 L of household ammonia (containing a minimum of 3 % ammonia) per 100 L of water while filling the tank with clean water. Agitate and then briefly flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full and allow to stand for 15 minutes with agitation. Flush the boom and hoses and drain tank completely.
4. Remove nozzles and screens and clean separately with an ammonia solution (100 mL in 10 L water).
5. Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume. Drain tank completely.

Note: If it is not possible to drain the tank completely, step 3 must be repeated before going onto step 4.

Caution: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which may cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

DIRECTIONS FOR USE

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEEDDED WITH LEGUMES.

2025-2245
2025-07-03

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat:
		Less than 1 m	Greater than 1 m	
Field sprayer	All labelled crops	1	0	3

When tank-mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

GORDEX™ A ALONE

Crops Registered

Spring wheat (including durum) and spring barley.

Application Directions

Apply 25 mL of **GORDEX™ A** per hectare in a minimum of 100 L per hectare of water. Add Agral 90 at 0.2% v/v. See weeds species controlled under “Weeds Controlled or Suppressed by

GORDEX™ A Alone at 25 mL/ha. Apply to actively growing wheat and barley from the 2 to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Application Timing

Apply to actively growing weeds in the 2-4 leaf stage. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Only weeds which are emerged at the time of application will be affected. If foliage is wet at the time of application control may be decreased. Under conditions of low crop and high weed density, control may be reduced.

DO NOT apply by air.

Weeds Controlled or Suppressed by GORDEX™ A Alone at 25 mL/ha

Weeds Controlled:

buckwheat, wild	shepherd's-purse
canola, volunteer♦	smartweed
chickweed, common	stinkweed
cleavers	mustard, wild

♦ Including herbicide-tolerant canola varieties except Clearfield® canola.

Weeds Suppressed:

hempnettle	sowthistle, annual
pigweed, redroot	sowthistle, perennial *

* Applications made at advanced leaf stages will reduce product effectiveness.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of **GORDEX™ A**
4. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
5. Add Agral® 90 as the last ingredient at the rate of 0.2% v/v (2 L per 1000 L of spray volume).
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

Preharvest/Grazing Intervals

Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.

TANK-MIX COMBINATIONS – GORDEX™ A + AXIAL® 100 EC HERBICIDE OR LADDER®

For control of annual grasses (see table below) tank mix **GORDEX™ A** with either Axial® 100 EC Herbicide or LADDER®. Refer to the above table for broadleaved weeds controlled or

2025-2245
2025-07-03

suppressed with **GORDEX™ A**.

Tank-Mix Combinations with GORDEX™ A for Annual Grass Control

Tank-Mix Partner	Crops Registered	Rate/ha	Adjuvant and Rate	Additional Weeds Controlled
Axial® 100 EC Herbicide	spring wheat spring barley	600 mL	Adigor® 700 mL/ha	wild oats, green foxtail♦ yellow foxtail, volunteer oats, volunteer canary seed, proso millet
LADDER®	spring wheat durum wheat	230 mL	XA Oil Concentrate or Score 0.8% v/v	wild oats, green foxtail, yellow foxtail, volunteer (tame) oats, volunteer canary seed, barnyard grass

♦Reduction in control of green foxtail may be observed when tank mixed with Axial 100 EC Herbicide.

♦Suppression only of green foxtail when tank mixed with Axial 100 EC Herbicide and MCPA Ester.

To control additional broadleaved weeds, MCPA Ester may be added to either tank mix up to a maximum of 420 g a.e./ha (equivalent to 840 mL/ha of MCPA LV500). Refer to the MCPA Ester tank mix partner label for rates less than 420 g a.e./ha and weed controlled.

Mixing Instructions

1. Begin to fill sprayer tank with clean water, and engage agitator. **Agitation must be continued throughout the entire mixing and spraying procedure.**
2. When the sprayer is three quarters full of water, add **GORDEX™ A** and agitate for 2-3 minutes.
3. If 2,4-D Ester, MCPA Ester or Curtail M® are also being added, add these next.
4. Add Axial® 100 EC Herbicide or LADDER®. Agitate for 2-3 minutes.
5. Add required Adjuvant.
6. Agitate for 1-2 minutes before adding remainder of water and then maintain constant agitation.
7. After any break in spraying operations, agitate thoroughly before spraying again.
8. **Use the spray suspension as soon as it is prepared.**
9. If an oil film starts to build up in the tank, drain tank and then clean as directed on this label.

TANK MIX COMBINATION - GORDEX™ A + MCPA Ester HERBICIDE

Crops Registered

Spring wheat (including durum), spring barley, and oats.

Application Directions

For control of a wide spectrum of broadleaved weeds apply **GORDEX™ A** at 25 mL/ha tank mixed with MCPA Ester Herbicide at 0.70 L/ha (350 g e.a./ha) in 100 L per hectare of water. Apply to actively growing wheat, barley, and oats from the 2 to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Weeds Controlled or Suppressed by the Tank Mixture of GORDEX™ A + MCPA Ester Herbicide

Weeds Controlled:

buckwheat, wild		ragweed, common
burdock♦♦	lamb's-quarters	shepherd's purse
canola, volunteer♦	mustard, ball	smartweed
chickweed, common	mustard, wild	stinkweed
cleavers	pigweed, redroot♦♦♦	sunflower, annual♦♦
flixweed (spring rosettes only)	pigweed, Russian♦♦	
hempnettle♦♦♦	prickly lettuce♦♦	

Weeds Suppressed:

dandelion (seedlings and overwintered rosettes less than 15 cm) ♦♦♦	sowthistle, perennial
plantain (top growth)	stork's bill♦♦♦
sowthistle, annual	thistle, Canada♦♦♦

♦Including herbicide-tolerant canola varieties

♦♦Up to 4 leaf stage

♦♦♦Optimum control can be achieved with the addition of 70 g a.e./ha) of MCPA Ester Herbicide

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of **GORDEX™ A**
4. Add the correct amount of MCPA Ester Herbicide and continue to agitate.
5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

Note: Do not add a surfactant to this tank mixture.

Pre-Harvest/Grazing Intervals (GORDEX™ A + MCPA Ester Herbicide)

1. Do not cut the treated crop for hay or graze treated crop within 7 days after application.
2. Do not harvest the treated crop within 60 days after application.

TANK-MIX COMBINATIONS - GORDEX™ A + MCPA Ester HERBICIDE + OTHER HERBICIDES

Crops Registered

Spring wheat (including durum) and spring barley.

Do not apply these tank mixes to oats.

Application Directions

Tank mixtures of **GORDEX™ A** + MCPA Ester Herbicide with other herbicides will provide control of broadleaved weeds and specified annual grasses. Apply to actively growing wheat and barley from the 2 to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Tank-Mix Combinations with GORDEX™ A + MCPA Ester Herbicide + Other Herbicides

Herbicide Tank-Mix Partner	Crops Registered	Rate (Product/ha)	Adjuvant and Rate	Additional Weeds Controlled
LADDER®	wheat (spring & durum)	230 mL	XA Oil Concentrate 0.8% v/v	wild oats, green foxtail, yellow foxtail, volunteer (tame) oats, volunteer canary seed, barnyard grass

♦Wheat (including durum) exposed to water-logged or saturated soils, or temperature extremes such as heat or freezing weather, or drought, low fertility or plant disease at application time could show unacceptable injury symptoms. Weed control may also be reduced by these same conditions.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add **GORDEX™ A**
4. Add MCPA Ester Herbicide and continue to agitate.
5. Add the required amount of tank mix partner and continue to agitate.
6. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
7. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
8. Follow sprayer clean-up directions.

Note 1: Add only the adjuvant recommended above. Follow tank-mix partner label for order of mixing and any specific instructions for mixing.

TANK MIX COMBINATION - GORDEX™ A + CURTAIL M HERBICIDE

Crops Registered

Spring wheat (including durum), spring barley, and oats.

Application Directions

For control of a wide spectrum of annual broadleaved weeds plus Canada thistle and perennial sowthistle, apply **GORDEX™ A** at 25 mL/ha tank mixed with 1.5 L/ha of Curtail® M Herbicide in 100 L per hectare of water. Apply to actively growing wheat, barley, and oats from the 2 to 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Weeds Controlled or Suppressed by the Tank Mixture of GORDEX™ A + Curtail® M Herbicide

Weeds Controlled:

buckwheat, wild	lamb's-quarters	stinkweed
canola, volunteer♦	mustard, wild	stork's-bill
chickweed, common	pigweed, redroot	thistle, Canada

2025-2245
2025-07-03

cleavers
dandelions (seedlings and overwintered rosettes < 15 cm)
flixweed (spring rosettes only)♦♦
sowthistle, annual

shepherd's purse
smartweed
hempnettle
sowthistle, perennial (top growth only) ♦♦

♦Including herbicide-tolerant canola varieties.

♦♦Control not observed until a minimum of 40 days after treatment.

Weeds Suppressed:

dandelion (overwintered rosettes > 15 cm, mature plants)

plantain (top growth)

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of **GORDEX™ A**
4. Add the correct amount of Curtail® M Herbicide and continue to agitate.
5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

Note: Do not add a surfactant to this tank mixture.

Pre-Harvest/Grazing Intervals (GORDEX™ A + Curtail® M Herbicide)

1. Do not cut the treated crop for hay or graze treated crop within 7 days after application.
2. Do not harvest the treated crop within 60 days after application.

TANK-MIX COMBINATION - GORDEX™ A + 2,4-D Ester HERBICIDE

Crops Registered

Spring wheat (including durum).

Application Directions

For control of a wide spectrum of broadleaved weeds apply **GORDEX™ A** at 25 mL/ha tank-mixed with 2,4-D Ester Herbicide at 560 g a.e./ha. Follow all precautions, directions for use and limitations on the tank -mix partner labels. Apply when wheat is in the 3 leaf expanded stage to the 6 leaf stage. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be affected. Best results are obtained from application made to seedling weeds.

Weeds Controlled or Suppressed by the Tank Mixture of GORDEX™ A + 2,4-D Ester Herbicide

Weeds Controlled:

bluebur	flixweed	ragweed, common
burdock	kochia♦♦	shepherd's purse
buckwheat, tartary	lady's-thumb	smartweed
buckwheat, wild	lamb's quarters	sowthistle, annual
canola, volunteer♦	mustard, ball	stinkweed
chickweed, common	mustard, wild	sunflower, annual

2025-2245
2025-07-03

cleavers	pigweed, redroot	thistle, Russian
cocklebur	plantain	vetch
dandelion (seedlings and overwintered rosettes)	prickly lettuce	wild radish

◆Including all herbicide-tolerant canola varieties.

◆◆Up to 5 cm in height

Weeds Suppressed:

Hempnettle,
sowthistle, perennial (top growth control),
thistle, Canada (top growth control)

Mixing Instructions

1. Fill sprayer tank $\frac{1}{2}$ full of water.
2. Start sprayer tank agitation.
3. Add **GORDEX™ A**
4. Add 2,4-D Ester Herbicide and continue to agitate.
5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

Pre-Harvest/Grazing Intervals (GORDEX™ A + 2,4-D Ester Herbicide)

1. Do not permit lactating dairy animals to graze fields within 7 days after application.
2. Do not harvest forage or cut hay within 30 days after application.
3. Withdraw meat animals from treated fields at least 3 days before slaughter.
4. Do not harvest the treated mature crop within 60 days after application.

TANK MIX COMBINATION WITH GORDEX™ A + BROMOXYNIL

Crops Registered

Spring wheat (including durum) and spring barley.

Application Directions

For control of a wide spectrum of broadleaved weeds, apply **GORDEX™ A** at 20-25 mL/ha tank mixed with 230 – 280 g e.a./ha of bromoxynil liquid formulations, including emulsifiable suspensions (for example, 0.96 – 1.2 L Bromotril® 240 EC, 0.8 – 1 L/ha Pardner® Herbicide or 0.96 – 1.2 L/ha Koril® 235 Liquid Herbicide), in 100 L per hectare of water. Apply to spring wheat, durum wheat, and spring barley from the 2 to 6 leaf stage. Apply when weeds are actively growing (2-4 leaf stage). Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Use the lower rate when the weeds infestations are low and the higher rate when the weeds infestations are heavy.

Weeds Controlled or Suppressed by GORDEX™ A + bromoxynil

Weeds Controlled: Up to 4 leaf stage

bluebur	kochia◆◆◆	smartweed, green
---------	-----------	------------------

2025-2245
2025-07-03

buckwheat, wild	lady's-thumb	smartweed, pale
canola, volunteer♦	mustard, wild	stinkweed
chickweed, common	nightshade, American	thistle, Russian♦♦
cleavers	pigweed, red-root	velvetleaf♦♦♦
cocklebur	ragweed, common	
cow cockle	shepherd's purse	

♦Including herbicide-tolerant canola varieties except Clearfield® canola.

♦♦Apply before plants are 5 cm high

♦♦♦Apply before plants are 8 cm high

Up to 8 leaf stage

buckwheat, common	groundsel, common
buckwheat, Tartary	lamb's-quarters

Weeds Suppressed:

hempenettle	sowthistle, perennial♦
sowthistle, annual	

♦Applications made at advanced leaf stages will reduce product effectiveness.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of **GORDEX™ A** and continue to agitate.
4. Add the correct amount of bromoxynil product
5. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

Note: Do not add a surfactant to this tank mixture.

Pre-Harvest/Grazing Intervals (GORDEX™ A + bromoxynil)

1. Do not cut the treated crop for hay or graze treated crop within 30 days of application.
2. Do not harvest the treated crop within 60 days of application.

TANK-MIX COMBINATIONS WITH GORDEX™ A + BROMOXYNIL + OTHER HERBICIDES

Crops Registered

Spring wheat (including durum).

Application Directions

Tank mixtures of **GORDEX™ A** + bromoxynil with other herbicides will provide control of broadleaved weeds and specified annual grasses. Apply when crops and weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Tank-Mix Combinations with GORDEX™ A + Bromoxynil Liquid Formulations, Including Emulsifiable Suspensions, + Other Herbicides

Tank-Mix Partner	Crops Registered	Rate/ha	Adjuvant and Rate	Additional Weeds Controlled
LADDER®	wheat (spring & durum)	230 mL	XA Oil Concentrate 0.8% v/v	wild oat and green foxtail

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add **GORDEX™ A**
4. Add bromoxynil product and continue to agitate.
5. Add tank mix partner and continue to agitate.
6. Add adjuvant if required and continue to agitate.
7. Fill sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
8. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
9. Follow sprayer clean-up directions.

TANK MIX COMBINATION - GORDEX™ A + GLYPHOSATE PRESENT AS ISOPROPYLAMINE SALT OR DIMETHYLAMINE SALT OR POTASSIUM SALT

Crops Registered

Spring barley, Oats, Spring wheat (including durum).

Application Directions

GORDEX™ A may be tank mixed with glyphosate present as isopropylamine salt or dimethylamine salt or potassium salt to control volunteer glyphosate-tolerant canola, wild buckwheat (up to 5 leaf) and top growth control of dandelion (up to 6 leaf stage), in addition to the labelled weeds controlled by **GORDEX™ A** or glyphosate.

Apply **GORDEX™ A** at 25 mL/ha with glyphosate (present as isopropylamine salt or dimethylamine salt or potassium salt) at 450 g a.e./ha in 100 L water per hectare.

Application Timing and Cropping

Apply to actively growing weeds in the 2-4 leaf stage, except where noted above. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control. Only weeds which are emerged at the time of application will be affected. If foliage is wet at the time of application, control may be decreased. Under conditions of high weed density, control, may be reduced.

Spring

GORDEX™ A + glyphosate may be applied in the spring prior to seeding and no longer than 48 hours after seeding prior to any crop emergence. Fields treated with **GORDEX™ A** in the spring may be planted to barley, oats, wheat, durum or summer fallowed.

Summer

Prior to August 1st: **GORDEX™ A** + glyphosate may be applied to summer fallow fields and seeded in the following spring to barley, canola, oats, peas or wheat (including durum) or summer fallowed.

After August 1st, **GORDEX™ A** + glyphosate may be applied to summer fallow fields and seeded in the following spring to barley, oats or wheat (including durum) or summer fallowed.

Fall

GORDEX™ A + glyphosate may be applied to stubble or summer fallow fields after August 1st and prior to freeze-up and may be seeded to barley, oats or wheat (including durum) or summer fallowed.

Do not use in successive years at the same site.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, **GORDEX™ A** is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to **GORDEX™ A** and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **GORDEX™ A** or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or integrated weed management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact ADAMA Agricultural Solutions

2025-2245
2025-07-03

Canada Ltd. at 1-855-264-6262.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label.

Gordex, Bromotril and Ladder are trademarks of an ADAMA Group Company.
All other products listed are registered trademarks of their respective companies.