



# OUTSHINE ALL IN<sup>®</sup>

A selective postemergence herbicide for the control of hard-to-kill annual broadleaf weeds in spring wheat, spring barley and oats. Now in a more convenient formulation!



## GROUP 2 & 4

### Active Ingredients:

Florasulam 2.5 g/L  
Fluroxypyr 100 g/L  
MCPA 350 g/L = EC

### Application Rates:

Rate: 405 ml/ac

### Packaging and Acres Treated:

Case: 2 x 8.1 L jugs  
Drum: 97.2 L drums  
Tote: 518 L

Acres Treated: 20 ac/jug;  
240 ac/drum;  
1280 ac/tote

### Water Volume:

Ground: 40 L/ac  
(10 US gal/ac)

### Rainfastness:

2 hours

### REGISTERED CROPS:

- Oats
- Spring Barley
- Wheat (spring, durum, winter)

### WEEDS CONTROLLED:

When applied between 2-4 leaf stage unless otherwise listed.

- |                         |                           |
|-------------------------|---------------------------|
| • Burdock               | • Smartweed               |
| • Cleavers (1-8 Whorl)* | • Stinkweed               |
| • Cocklebur             | • Storksbill <sup>s</sup> |
| • Common Chickweed      | • Sunflower (annual)      |
| • Flixweed              | • Vetch                   |
| • Hemp-Nettle           | • Volunteer Canola        |
| • Kochia*               | • Volunteer Flax          |
| • Lamb's Quarters       | • Wild Buckwheat          |
| • Plantain              | • Wild Mustard            |
| • Prickly Lettuce       | • Wild Radish             |
| • Ragweed               |                           |
| • Redroot Pigweed       |                           |
| • Russian Pigweed       |                           |
| • Shepherd's Purse      |                           |

\* Including ALS resistant biotypes | <sup>s</sup> Suppression

### HOW IT WORKS:

OUTSHINE ALL IN<sup>®</sup>, applied early and thoroughly to the main flush of actively growing broadleaf weeds, quickly causes plants to stop growing, even if typical symptoms of dying weeds are not noticeable for 1–2 weeks after application.

# OUTSHINE ALL IN®

## CROP STAGING:

2-leaf expanded to just prior to flag leaf emergence.

## REGISTERED AND SUPPORTED TANK MIXES:

### Barley:

- Assert® 300 SC
- BRAZEN® II
- BRAZEN ALL IN®

### Wheat (spring, durum):

- |                  |                  |                     |
|------------------|------------------|---------------------|
| • Assert® 300 SC | • CAZADO™        | • Simplicity® GoDRI |
| • BRAZEN® II     | • Everest® 3.0   | • Traxos®           |
| • BRAZEN ALL IN® | • LADDER ALL IN® | • Varro®            |

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

## MIXING INSTRUCTIONS:

1. Fill sprayer tank ½ full of water.
2. Start sprayer tank agitation.
3. Add the WG herbicides first, then liquids, depending on the mixture followed by OUTSHINE ALL IN® and continue to agitate.
4. Add the required amount of tank-mix partner.
5. Fill the sprayer tank with sufficient water to spray 10 US gal/ac.

NOTE: Add only the adjuvant recommended. Follow tank-mix partner label for order of mixing. If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

## CROP ROTATIONS:

Fields previously treated with OUTSHINE ALL IN® can be seeded the following year to alfalfa, barley, canola, corn, fababeans, field beans, flax, lentils, mustard, oats, peas, potato, soybean, sunflower or wheat or fields can be summerfallowed.

## GRAZING RESTRICTIONS:

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 7 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

## PRE-HARVEST INTERVAL:

Do not harvest the treated crop within 60 days after application.

## STORAGE:

- Do not freeze.

[CLICK HERE FOR FULL  
PRODUCT DETAILS.](#)



## QUICK TIPS:

2 unique modes of action and 3 active ingredients provide resistance management and high performance.



Always read and follow label directions.  
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

BRAZEN®, BRAZEN ALL IN®, LADDER ALL IN® and OUTSHINE ALL IN® are registered trademarks and CAZADO™ is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies.  
© 2026 ADAMA Agricultural Solutions Canada Ltd.