



ADAMA

# VANTANA™

This Group 29 fungicide is an important resistance management tool for dry beans, potatoes and soybeans.



GROUP 29

## Active Ingredient:

Fluazinam 500 g/L = SC

## Application Rates:

Rate: 162–906 ml/ac

## Packaging

## & Acres Treated:

Case: 2 x 10 L jugs

Acres Treated: 11–62 ac/jug

## Water Volume:

Ground: Varies with the crop.

Refer to the label for more details.

Aerial: Minimum of 18.2 L/ac

## Rainfastness:

Avoid applying when heavy rainfall is forecast.

## KEY BENEFITS:

- Resistance Management Tool
- Can be used on multiple crops
- Broad spectrum disease control
- Multiple rates for precise control

## REGISTERED AND SUPPORTED CROPS:

- Dry-Shelled Beans
- Potatoes
- Soybeans

For a full list of registered crops, please refer to the label.

## HOW IT WORKS:

VANTANA™ works by inhibiting fungal adenosine triphosphate (ATP) production in the mitochondria (impairing energy production).

Active on a wide range of diseases, VANTANA™ is a protective fungicide that inhibits the germination of fungal spores.

## MIXING INSTRUCTIONS:

1. Fill sprayer tank ½ full of water.
2. Start sprayer tank agitation.
3. Add the required amount of VANTANA™.
4. If applicable, add any required adjuvants or surfactants for tank-mix partners.
5. Complete filling tank with water.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

FOR FULL  
PRODUCT  
DETAILS,  
SCAN THE  
QR CODE.



# VANTANA™

## APPLICATION TIMING AND KEY DISEASES CONTROLLED BY CROP:

NOTE: For application timing by crop, please refer to the label.

Crop	Disease	Rate
Dry-Shelled Beans	White Mould	243–405 ml/ac
	Anthracnose	
Potatoes	Late Blight	162–242 ml/ac
	White Mould	356–473 ml/ac
Soybeans	White Mould <sup>s</sup>	178 ml/ac
	White Mould	356–473 ml/ac

<sup>s</sup>Suppression

## CROP ROTATIONS:

Areas treated with VANTANA™ may be replanted with potatoes and dry shelled beans as soon as practical after the last application. Other root crops and leafy vegetables can be planted 30 days after the last application. All other crops can be planted 70 days after the last application.

## GRAZING RESTRICTIONS:

Do not feed treated foliage to livestock.

## RE-ENTRY INTERVALS

All other crops: 24 hours

## PRE-HARVEST INTERVALS:

- Dry-Shelled Beans: 30 days
- Potatoes: 14 days
- Soybeans: DO NOT apply after growth stage R3

## REGISTERED AND SUPPORTED TANK MIXES:

None registered.

## STORAGE:

Do not freeze.



Always read and follow label directions.  
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

VANTANA™ is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.

## QUICK TIPS:

For optimal efficacy VANTANA™ requires good coverage, best achieved with higher water volumes. Actual water volumes will vary with the crop, please refer to the label for more specific details.