



ADAMA

RUSH 24 ALL IN[®]

Controls a wide spectrum of broadleaf weeds like Cleavers and Kochia (including Group 2-resistant biotypes) with excellent grassy weed herbicide compatibility. Now in a more convenient formulation!



GROUP 4

Active Ingredients:

Fluroxypyr 90 g/L and 2,4-D Ester 360 g/L as an EC

Application Rates:

Rate: 445 ml/ac

Packaging and Acres Treated:

Case: 2 x 8.9 L jug

Drum: 106.8 L

Tote: 427.2 L

Acres Treated: 40 ac/case;
240 ac/drum;
960 ac/tote

Water Volume:

Ground: 12-40 L/ac
(3-10 US gal/ac)

Aerial: 12-20 L/ac
(3-5 US gal/ac)

Rainfastness:

1 hour

KEY BENEFITS:

- Control of Group 2-resistant biotypes of Kochia
- Suppression of Group 2-resistant biotypes of Chickweed
- Convenient and trusted formulation
- Easily tankmixes with the most popular grassy weed control herbicides
- Available in convenient bulk options

REGISTERED CROPS:

- Barley
- Wheat (durum, spring, winter)

WEEDS CONTROLLED:

Apply postemergence when weeds are in the seedling stage (2-4 leaf, unless otherwise noted)

- | | | |
|--------------------------------|---|-----------------------------|
| • Bluebur | • Kochia ² | • Stinkweed |
| • Burdock | • Lamb's Quarters | • Sunflower (annual) |
| • Cleavers (1-8 whorls) | • Mustards (except green tansy, dog & grey tansy) | • Vetch |
| • Clovers (sweet) | • Plantain | • Volunteer Canola |
| • Cocklebur | • Prickly Lettuce | • Volunteer Flax (1-12 cm) |
| • Flixweed | • Ragweed | • Wild Buckwheat (1-6 leaf) |
| • Field Horsetail ¹ | • Round-Leaved Mallow (1-6 leaf) | • Wild Radish |
| • Goat's-Beard | • Shepherd's Purse | |
| • Hemp-Nettle (2-6 leaf stage) | • Stork's Bill (1-8 leaf) | |
| • Hoary Cress ¹ | | |

WEEDS SUPPRESSED:

- | | | |
|--|-------------------|--|
| • Common Chickweed ² (up to 3 inches) | • Redroot Pigweed | • Sow Thistle (perennial) ¹ |
|--|-------------------|--|

Additional weeds controlled/suppressed with addition of 81 ml/ac (2 oz/ac) of 2,4-D ESTER 700:

- | | | |
|-------------------------------|-----------------------------|---|
| • Blue Lettuce ¹ | • Hairy Galinsoga | • Russian Thistle |
| • Dandelion (spring rosettes) | • Hedge Bindweed | • Smartweed |
| • Docks | • Lady's Thumb | • Tartary Buckwheat |
| • Field Bindweed ¹ | • Leafy Spurge ¹ | • Common Chickweed ² (up to 8 cm) |
| • Field Peppergrass | • Mustard (dog, tansy) | • Canada Thistle ¹ |
| • Gumweed | • Oak-Leaved Goosefoot | • Sow Thistle (perennial ¹ , annual) |
| | • Redroot Pigweed | |

¹Top growth control only

²Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme.



CLICK HERE FOR FULL PRODUCT DETAILS.



RUSH 24 ALL IN®

HOW IT WORKS:

Group 4 herbicides disrupt normal plant growth, resulting in twisting and cupping of leaves, epinasty and death of susceptible plants in 2–10 days.

CROP STAGING:

- 4-leaf to just prior to flag leaf emergence.
- Apply to Winter Wheat in the spring from the 3 tiller stage to just before the flag leaf stage.

REGISTERED AND SUPPORTED TANK MIXES:

Wheat only:

- CAZADO™
- Everest® 3.0
- LADDER ALL IN®
- Simplicity™ GoDRI™¹
- Traxos®
- Varro®

Wheat and Barley:

- Assert® 300 SC
- BISON® 400 L
- Puma® Advance

¹ Additional 2,4-D ester is not recommended when mixing RUSH 24 ALL IN® and Simplicity™.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

1. Fill the clean spray tank ½ full of clean water.
2. Add the required amount of RUSH 24 ALL IN™ and agitate thoroughly, followed by the required amount of additional 2,4-D ESTER 700 herbicide, if required.
3. Continue agitation and add the required amount of the tank-mix partner.
4. Complete filling the tank to the desired level with water.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

CROP ROTATIONS:

The following crops may be grown one (1) year after application:

- Barley
- Canola
- Flax
- Forage Grass
- Lentils
- Mustard
- Oats
- Field Peas
- Rye
- Wheat

GRAZING RESTRICTIONS:

- Do not permit lactating dairy animals to graze fields within seven (7) days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least three (3) days before slaughter.

PRE-HARVEST INTERVAL:

Barley, Wheat (durum, spring): 60 days

STORAGE:

Do not freeze.

QUICK TIPS:

RUSH 24 ALL IN® activity is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, for example drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

BISON®, LADDER ALL IN® and RUSH 24 ALL IN® are registered trademarks and CAZADO™ is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.