



CAPTAN® 80 WSP

A water-dispersible granule sealed within individual water-soluble pouches. The pouches and pesticide will dissolve readily in water for use as a spray for the control of certain fungus diseases of fruit, vegetables and ornamental crops. CAPTAN® 80 WSP fungicide is also useful as a soil treatment for the control of certain seed-rots and damping-off diseases.



Active Ingredient:

Captan

Fungicide Group:

Group M4

Formulation:

80% CAPTAN = WSP
(water-soluble pouch)

Application Rates and Packaging:

- Rates vary by crop, see label
- 4 foil bags containing 5 pouches of 0.5 kg per case

REGISTERED CROPS:

- Apples
- Apricots
- Cherries
- Cucumbers (field grown)
- Ginseng
- Grapes
- Highbush blueberries, blackberries, loganberries
- Lowbush blueberries
- Nectarine trees
- Peaches
- Pears
- Plums
- Potatoes
- Prunes
- Raspberries
- Rhubarb
- Strawberry (field grown)
- Tomatoes (field grown – foliar applications)
- Turf (golf course, sod farms only)
- Outdoor ornamentals (cut and non-cut flowers)
- Greenhouse ornamentals (non-cut flower: camellia, carnation, chrysanthemum, dahlia, foliar applications to aster, lilac, rose and tulip)
- Soil and greenhouse bench treatment (soil treatment: beans, celery, crucifers, eggplants, lawn seedbeds, peas, peppers, seedlings or transplants of roses and other flowers, shrubs, tomatoes, trees)

KEY DISEASES CONTROLLED AND APPLICATION RATES:

Crop	Disease	Application Rate
FRUIT CROPS		
Apples	Scab, sooty blotch, fly speck, Brook's spot, bitter rot, black rot, bull's eye rot	1.2 kg/ac
Pears	Scab, sooty blotch	
Apricots	Brown rot	1.6 kg/ac
Cherries	Brown rot, leaf spot (Shot Hole)	
Peaches and Nectarines	Brown rot, scab	
Plums and Prunes	Black knot, brown rot	
Grapes	Dead arm (current season's infections)	0.8 kg/acre
	Downy mildew, black rot	0.8–1.2 kg/acre
Raspberries	Fruit rot spur blight	1.0 kg/400 L of water per acre
Blackberries	Fruit rot	0.9 kg/400 L of water per acre
Loganberries	Cane spot, fruit rot, leaf spot, spur blight	0.6–0.9 kg/400 L of water per acre



CAPTAN® 80 WSP

Crop	Disease	Application Rate
FRUIT CROPS (continued)		
Blueberries	Fruit rot, mummy berry	0.9 kg/400 L of water per acre
Strawberries	Grey mould, leaf spot	1.4 kg/acre
Rhubarb (in forcing sheds)	Leaf rot	0.5–0.8 kg/acre
VEGETABLE CROPS		
Cucumbers (field only)	Anthrachnose, scab	0.9–1.7 kg/acre
Potatoes	Early blight, late blight	1.0–1.5 kg/acre
Tomatoes	Anthrachnose, Septoria leaf spot	1.1–1.7 kg/acre
Ginseng	Rhizoctonia root rot, pythium root rot and damping-off, phytophthora root rot and grey mould, cylindrocarpon root rot ¹	1.0 kg/ac (0.8 kg ai/ac) in a spray volume of 378–757 L of water/acre

¹Suppression

WATER VOLUME:

- Ground: 400 L/ac
- Aerial: 400 L/ac (some restrictions, see label)

REGISTERED AND SUPPORTED TANK MIXES:

Nova™

MIXING INSTRUCTIONS:

1. Fill the spray tank $\frac{1}{3}$ to $\frac{1}{2}$ full with clean water and begin agitation or bypass.
2. Add the required number of unopened pouches of CAPTAN® 80 WSP fungicide directly to the spray tank.
3. Vigorous agitation is required for CAPTAN® 80 WSP fungicide to become fully suspended. The water temperature and the degree of agitation will determine the amount of time for the pouches to dissolve. The pouches should be completely dissolved before application or adding tank-mix partners. Maintain sufficient agitation during both mixing and application.
4. If required, add the tank-mix partner.

CROP ROTATIONS:

No restrictions

GRAZING RESTRICTIONS:

N/A

STORAGE:

No restrictions

PRE-HARVEST INTERVAL:

- Apricots, blueberries, cherries, cucumbers, loganberries, peaches, nectarines, potatoes, plums, prunes, rhubarb, strawberries, tomatoes: 2 days
- Apples, pears, grapes: 7 days
- Ginseng: 20 days

QUICK TIPS:

The pouches containing CAPTAN 80 WSP fungicide are water-soluble and will dissolve completely in water. After opening the outer bag, drop the required number of unopened inner pouches into the spray tank as directed. Reseal outer bag to protect remaining pouches. Do not excessively handle water-soluble pouches or expose to moisture since this may cause breakage. Do not allow pouches to become wet prior to mixing spray solution.



Always read and follow registered product label instructions. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

Toll-free: 1.855.264.6262
Website: ADAMA.COM

®/™ CAPTAN is a registered trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies.
© 2021 ADAMA Agricultural Solutions Canada Ltd.