




ADAMA

CUSTODIA®

A multi-mode of action fungicide offering preventative and curative protection of the flag leaf against all major leaf diseases in wheat and barley.



Active Ingredient:

Tebuconazole and Azoxystrobin

Fungicide Groups:

Group 3 and Group 11

Formulation:

Tebuconazole 200 g/L and Azoxystrobin 120 g/L = SC

Application Rates and Packaging:

- 0.19 L/ac – 0.25 L/ac
- 400 – 525 acres/jug
- 2 x 10.08 L case

Registered crops:

Wheat (spring, winter), Barley

KEY BENEFITS:

- Multiple modes of action for resistance management
- Curative and preventative disease control
- Broad-spectrum leaf disease control with superior rust control compared to competitors
- Penetrates quickly into the leaf tissue
- Keeps the plant greener longer

KEY DISEASES CONTROLLED:




- Leaf rust
- Stem rust
- Stripe rust
- Septoria leaf blotch
- Tan spot
- Net blotch
- Spot blotch

HOW IT WORKS:

CUSTODIA® is a broad-spectrum fungicide with both preventative and curative properties. With its unique systemic and translaminar activity, CUSTODIA® has shown to both move through the leaves and down the plant for more protection. CUSTODIA® penetrates quickly into the leaf tissue protecting the flag leaf from all major leaf diseases.

CUSTODIA®

Field Test Results



NOTE: Means followed by same letter do not significantly differ (P=.05, Student-Newman-Keuls)


CUSTODIA™

CROP STAGING:

Crop	Diseases	Application Timing	Rate
Wheat (spring, winter)	Leaf rust Stem rust Stripe rust Septoria leaf blotch Tan spot	Apply CUSTODIA™ to leaf foliage at the first sign or very early stage of disease, up to the beginning of heading. Use of the higher rate should be considered when weather conditions are conducive to heavy disease development.	0.19 L/ac – 0.25 L/ac
Barley	Net blotch Spot blotch Leaf rust Stem rust Stripe rust Septoria leaf blotch Tan spot	Apply CUSTODIA™ to leaf foliage at the first sign or very early stage of disease, up to the beginning of heading. Use of the higher rate should be considered when weather conditions are conducive to heavy disease development.	0.19 L/ac – 0.25 L/ac

Orchestrated Leaf Disease Control

Made just for cereals, our portfolio of fungicides are formulated to help you manage some of the most damaging leaf diseases. Use this guide to help you determine the best product and spray timing for your crop, for disease protection when you need it most.





CUSTODIA®

WATER VOLUME:

Ground: 40 L/ac or 10 gal/ac

Air: 20 L/ac or 5 gal/ac

RAINFASTNESS:

Avoid application when heavy rain is forecast.

MIXING INSTRUCTIONS:

1. For ground application use 40 L/ac, for aerial use 19 L/ac.
2. Use a 50-mesh (or coarser) filter screen.
3. Fill the spray tank $\frac{3}{4}$ full with water.
4. Add the required amount of CUSTODIA® foliar fungicide into the sprayer.
5. Agitate until the fungicide is thoroughly mixed.
6. Continue agitation and add the required amount of the tank-mix partner.
7. Complete filling the tank to the desired level with water.
8. Upon completion of spraying, thoroughly flush tank, boom, hoses and in-line and nozzle screens with clean water to avoid possible injury to other crops.
9. Repeat sprayer cleanout process using an appropriate spray system cleaner.

CROP ROTATIONS:

No restrictions

PRE-HARVEST INTERVALS:

Mature grain: 36 days

Forage and hay: 6 days

GRAZING RESTRICTIONS:

Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment. Straw cut after harvest may be fed or used for bedding.

STORAGE:

1. Store above 5°C in original, tightly closed container.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. Keep from freezing.

Quick Tips

CUSTODIA® should be applied at flag leaf for optimal leaf disease control.

Coverage is key; do not use less than recommended water volumes.

For fusarium control, we would recommend an application of ORIUS® 430 SC fungicide at full head emergence after CUSTODIA® at flag leaf.

For resistance management, CUSTODIA® contains Group 3 and 11 fungicides.

When possible, rotate the use of CUSTODIA® or other Group 3 and 11 fungicides with different groups that control the same pathogens.



ADAMA

Always read and follow label directions.
Toll-free: 1.855.264.6262
Website: ADAMA.COM/CANADA

®/™ BUMPER, CUSTODIA and ORIUS are registered trademarks of ADAMA Agricultural Solutions Canada Ltd.
All other products are trademarks of their respective companies. © 2021 ADAMA Agricultural Solutions Canada Ltd.