



ESTEEM®

Superior control of hard to control broadleaf weeds like thistle, cleavers and kochia in barley, spring wheat and durum.



Herbicide Group

Group 4

Active Ingredients:

Fluroxypyr 180 g a.e./L,
MCPA 600 g a.e./L = EC and
Clopyralid 360 g a.e./L = SL

Application Rates and Acres Treated:

Rate:

Low: 240 ml/ac Fluroxypyr;
280 ml/ca MCPA Ester 600;
84 ml/ac Clopyralid

High: 320 ml/ac Fluroxypyr;
365 ml/ac MCPA Ester 600;
111 ml/ac Clopyralid

Acres Treated: 30 – 40 ac/case

Packaging:

Case: Fluroxypyr 180: 1 × 9.6 L jug;
MCPA Ester 600: 1 × 11.01 L jug;
Clopyralid 360: 1 × 3.32 L jug

REGISTERED CROPS:

- Barley
- Wheat (spring, durum)

WEEDS CONTROLLED:

At the low rate of 40 ac/case will control:

Wild buckwheat, burdock, cleavers, Canada thistle (low infestations), cocklebur, field horsetail (top growth), volunteer flax, flixweed, kochia, lamb's quarters, wild mustard, plantain (top growth), prickly lettuce, ragweeds, shepherd's purse, stinkweed, stork's bill, volunteer sunflowers, annual sunflowers, vetch, wild radish

At the high rate of 30 ac/case will control:

Tartary buckwheat, Canada thistle (medium to high infestations, season long control), volunteer canola, chickweed, dandelions, common groundsel, hempnettle, roundleaf mallow, redroot pigweed, Russian pigweed, scentless chamomile, smartweed, annual sow thistle, perennial sow thistle (season long control)

ESTEEM®

HOW IT WORKS:

The components of ESTEEM® move within the plant to control exposed and underground plant tissues. It mimics naturally occurring plant hormones which control weeds by disrupting normal plant growth patterns. Symptoms of effect include epinasty (twisting of the stems) and swollen nodes.

CROP STAGING:

3-leaf stage to just before flag emergence

RAINFASTNESS:

4 hours

WATER VOLUME:

- Ground: 20 – 40 L/ac (5 – 10 US gal/ac)
- Aerial: 12 – 20 L/ac (3 – 5 US gal/ac)

REGISTERED AND SUPPORTED TANK MIXES:

- Assert® 300 SC
- BISON® 400 L
- BRAZEN™ II
- BroadBand®
- LADDER ALL IN®
- Puma® Advance
- Traxos®
- Varro®

For tank mixes with registered pest control products, the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Buffer Zones must be followed for each product. In cases where these requirements differ between the tank-mix partner labels, the most restrictive label must be followed.

MIXING INSTRUCTIONS:

1. ½ fill the tank with clean water.
2. Add the required amount of ADAMA clopyralid, agitate thoroughly.
3. Add the required amount of MCPA ester and fluroxypyr, agitate thoroughly.
4. Add any tank-mix partners, agitate thoroughly.
5. Fill the tank and agitate again before using.

CROP ROTATIONS:

Barley, canola, flax, forage grasses, mustard, oats, rye and wheat can be seeded the following year. If rotating to peas or soybeans, a rainfall of 140 mm (5.5 inches) between herbicide application and August 31, and an annual precipitation greater than 175 mm (6.9 inches) is required.

STORAGE:

Do not freeze.

PRE-HARVEST INTERVAL:

Do not harvest treated crop within 60 days of application.

GRAZING RESTRICTIONS:

Do not graze livestock within 7 days of application.



ADAMA

Always read and follow label directions.
Toll-free: 1.855.264.6262
Website: ADAMA.COM/CANADA

BISON®, ESTEEM® and LADDER ALL IN® are registered trademarks, and BRAZEN™ is a trademark of ADAMA Agricultural Solutions Canada Ltd. All other products are trademarks of their respective companies. © 2021 ADAMA Agricultural Solutions Canada Ltd.