

# **ZIVATA**<sup>™</sup>

New choice in insect control giving you the same trusted results in a more sustainable advanced formulation.





# Insecticide Group:

Group 3

# **Active Ingredient:**

Lambda-cyhalothrin 120g/L as an EC

# Application Rates and Acres Treated:

Rate: 17 – 94 ml/ac (standard rate for most pests: 34 ml/ac)

Acres Treated: 45 – 240 ac/jug (120 ac/jug at standard rate)

# Packaging:

Case: 2 × 4.08 L jugs

#### **KEY BENEFITS:**

- Advanced formulation using sustainable plant-based materials
- Low Volatile Organic Compound (VOC) with low odour formulation
- Improved user experience and flexibility of use around odour-sensitive areas
- Trusted and proven active ingredient
- Broad range of crops, pests and use patterns

## **REGISTERED CROPS:**

- Alfalfa
- Beans
- Canola
- Cereals (wheat, barley, oats)
- Corn (field)

- Flax
- Potatoes
- Soybeans
- Sunflowers

 $\mathsf{ZIVATA}^{\mathsf{M}}$  is registered for use on more than 30 crops; refer to the label for more information.

#### **HOW IT WORKS:**

ZIVATA™ is a synthetic pyrethroid insecticide formulated with an improved, plant-based solvent that offers fast-acting stomach and contact effects against a broad spectrum of insect pests. This renewably sourced formulation has low volatile organic properties and improves the user experience with a reduced drift potential and product volatility.



#### **WATER VOLUME:**

• Ground: 10 – 20 US gal/ac (40 – 80 L/ac)

• Aerial: 1-5 US gal/ac (4-16 L/ac)

### **RAINFASTNESS:**

1 hour

#### **REGISTERED AND SUPPORTED TANK MIXES:**

#### Herbicides:

- Assert®
- BISON® 400 L
- BRAZEN™ II
- Everest<sup>®</sup>
- LADDER ALL IN®
- SQUADRON® II

## Fungicides:

- Allegro<sup>®</sup>
- BUMPER® 432 EC
- SORATEL™
- TOPNOTCH™

## **MIXING INSTRUCTIONS:**

Compatibility should always be confirmed by premixing small proportional quantities of water, ZIVATA™, and the tank-mix partner in advance.

#### **CROP ROTATIONS:**

No restrictions.

### **PRE-HARVEST INTERVAL:**

- Corn (silage): 14 days
- · Corn (field): 21 days
- Legumes (soybeans, beans, field peas, faba beans, chickpeas, lentils): 21 days
- · Oilseeds: 7 days

- Potatoes: 7 daysSunflowers: 7 daysTimothy: 14 days
- Wheat, Barley, Oats: 28 daysWheat for forage: 14 days

## **GRAZING RESTRICTIONS:**

Do not graze livestock within 3 days of application in alfalfa and 14 days of application in wheat, barley, oats, and timothy.

For silage corn do not apply within 14 days of harvesting for silage. All other registered crops DO NOT graze, or harvest treated forage, straw, or hay for livestock feed.

# **STORAGE:**

Do not freeze.

# **QUICK TIPS:**

Control of some insect species with pyrethroid insecticides decreases as temperature rises (above 25° C). For best results, apply ZIVATA™ during the early morning before temperatures rise, and during the evening, past the heat of the day. Use sufficient water for thorough coverage.



Always read and follow label directions. Toll-free: 1.855.264.6262 Website: ADAMA.COM