

# **Active Ingredients:**

Fluroxypyr 180 g/L, Bromoxynil 225 g/L, MCPA 2 EH Ester 225 g/L = EC

# Application Rates and Acres Treated:

### Rate:

500 ml/ac Bromoxynil/MCPA Ester; 240 ml/ac Fluroxypyr

### Acres Treated:

40 ac/case; 480 ac/pallet

## Packaging:

### Case:

Bromoxynil/MCPA Ester: 2 × 10 L jugs; Fluroxypyr: 1 × 9.6 L jug

### Pallet:

Bromoxynil/MCPA Ester: 2 × 120 L drums; Fluroxypyr: 1 × 115.2 L drum

### **Water Volume:**

Ground: 20 – 40 L/ac (5 – 10 US gal/ac) Aerial: Do not apply.

## Rainfastness:

1 hour

# GROUP 4 & 6

# REGISTERED CROPS: • Barley

WEEDS CONTROLLED:

### • American nightshade

- Bluebur
- bluebu
- Burdock
- Canada thistle<sup>1</sup>
- Chickweed
- Cleavers (including Group 2 resistant biotypes)
- Cocklebur
- Common groundsel
- Cow cockle
- Flixweed
- Hempnettle
- Kochia (including Group 2 and glyphosate-resistant biotypes)
- · Lady's thumb

<sup>1</sup>Top growth control.

- Wheat (spring, durum)
- · Lamb's quarters
- Night-flowering catchfly
- · Perennial sow thistle
- Redroot pigweed
- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed
- Vetch
- Volunteer canola
- Volunteer sunflower
- Wild buckwheat
- Wild mustard (including Group 2 resistant biotypes)
- · Wild radish
- Wormseed mustard

### **HOW IT WORKS:**

Quickly causes plants to stop growing. Convenient co-pack controls a wide range of weeds, including glyphosate-resistant and Group 2 resistant kochia, Group 2 resistant cleavers and Group 2 resistant wild mustard.





# **FORCEFIGHTER® M**

### **CROP STAGING:**

Crop	Stage
Wheat	2 leaf to early flag leaf
Barley	2 leaf to early flag leaf

#### **REGISTERED AND SUPPORTED TANK MIXES:**

### Spring wheat:

- BISON® 400 L
- BRAZEN™ II
- Everest® 3.0
- LADDER ALL IN®
- Simplicity® GoDRI
- Refine® SG
- Traxos®

### Durum wheat:

- Everest® 3.0
- LADDER ALL IN®
- · Simplicity® GoDRI
- Traxos®

### **Barley:**

- BISON® 400 L
- BRAZEN™ II

#### MIXING INSTRUCTIONS:

- 1. Fill spray tank ½ full with water.
- 2. Add the required amount of FORCEFIGHTER® M and agitate thoroughly.
- 3. Add any tank-mix partners.
- 4. Fill the tank and agitate again before use.

### **CROP ROTATIONS:**

Can be seeded the following year to barley, canola, flax, forage grasses, lentils, mustard, field peas, rye, and wheat.

### **GRAZING RESTRICTIONS:**

30 days

### STORAGE:

Do not freeze.

### **PRE-HARVEST INTERVAL:**

60 days

### **QUICK TIPS:**

FORCEFIGHTER® M's activity is influenced by weather conditions. The optimal temperature is 12 – 24° C. Avoid application 3 days before or after frost. Do not apply before the 2-leaf stage as crop injury may occur. Use 38 L/ac application volume when there is a heavy canopy or when most weeds are at an advanced stage of growth. Do not apply by air.

