



ADAMA

GLUFOSINATE 150 SL

Reliable formulation for post-emergent control of almost 30 broadleaf and grassy weeds for use in glufosinate-ammonium-tolerant canola and soybeans.

GROUP 10

Active Ingredient

Glufosinate ammonium
150 g/L = SL

Packaging

Tote: 432 L

Application Rates & Acres Treated

Rate: 0.8 – 1.62 L/ac

Acres treated:
267 – 540 ac/tote

Water Volume

Ground: 45 L/ac (12 US gal/ac)

Aerial: 23 L/ac (6 US gal/ac)

Rainfastness

4 hours

REGISTERED CROPS

- Glufosinate-ammonium-tolerant canola
- Glufosinate-ammonium-tolerant soybeans

Crop	Timing	Recommended Rates
Canola	Apply from the cotyledon stage up until, but prior to, the early bolting stage of canola.	<p>One (1) pass: 1.35 – 1.62 L/ac</p> <p>Two (2) passes: 1.35 L/ac followed by 1.35 L/ac (and up to 1.62 L/ac)</p> <p>OR</p> <p>1.62 L/ac followed by 1.35 L/ac</p> <p>NOTE: Do not apply more than a total of 2.97 L/ac in one season.</p>
Soybeans	Apply from the cotyledon to the flowering stage of the crop.	0.8 – 1.35 L/ac

NOTE: Please refer to label for additional rates.

HOW IT WORKS

ADAMA GLUFOSINATE 150 SL is a non-selective herbicide that provides control of a broad spectrum of grassy and broadleaf weeds in canola and soybean varieties and hybrids that are specially developed to be tolerant to glufosinate ammonium.

ADAMA GLUFOSINATE 150 SL is a contact herbicide with limited translocation within the plant. Control is best when weeds are actively growing and not under stress.



ADAMA GLUFOSINATE 150 SL

WEEDS CONTROLLED

BROADLEAF WEEDS

- Canada thistle¹
- Cleavers
- Common chickweed
- Cow cockle
- Dandelion
- Flixweed
- Hemp-nettle
- Kochia
- Lady's thumb
- Lamb's quarters
- Perennial sow thistle
- Redroot pigweed
- Round-leaved mallow
- Russian thistle
- Scentless chamomile
- Shepherd's purse
- Smartweed
- Stinkweed
- Stork's bill
- Volunteer flax
- Wild buckwheat
- Wild mustard

GRASSES

- Barnyard grass
- Green foxtail
- Quackgrass³
- Volunteer barley²
- Volunteer wheat
- Wild oats

¹ Top growth suppression only

² Suppression only

³ Season-long control for heavy populations at 1.62 L/ac

REGISTERED AND SUPPORTED TANK MIXES

Herbicides:

- Facet[®] L
- ARROW ALL IN[®]
- LEOPARD[®]

MIXING INSTRUCTIONS

1. Fill the tank $\frac{3}{4}$ full with clean water.
2. Add the correct amount of ADAMA GLUFOSINATE 150 SL.
3. Add the remaining amount of water, begin agitation, and spray out immediately.

CROP ROTATIONS

Buckwheat, barley, millet, oats, rye, sorghum, triticale: 70 days

All other crops (except field corn, canola and soybeans, dry common beans (not grown for seed), alfalfa, carrots, lettuce, onion and potatoes): 120 days

PRE-HARVEST INTERVAL

Canola: 60 days

Soybeans: 70 days

STORAGE

Do not freeze.

GRAZING RESTRICTIONS

Canola:

- Grain and meal from treated crop can be fed to livestock.
- Do not graze the treated crop or cut for hay; sufficient data is not available to support such use.

Soybeans: 20 days

QUICK TIPS:

ADAMA GLUFOSINATE 150 SL can be weak on some annual grasses. If these species are not adequately controlled with a residual herbicide, it may be advantageous to include a grass herbicide such as LEOPARD[®] or ARROW ALL IN[®] with the post-emergent glufosinate application.

ADAMA GLUFOSINATE 150 SL activity is maximized in warm, sunny weather.

ADAMA GLUFOSINATE 150 SL, as with all ADAMA products, is fully supported by a sales, service and agronomic teams across all Western Canadian provinces.



ADAMA

Always read and follow label directions.
1.855.264.6262 ADAMA.COM

^{®/™} ARROW ALL IN and LEOPARD are registered trademarks of ADAMA Agricultural Solutions Canada Ltd. All products are trademarks of their respective companies.

© 2023 ADAMA Agricultural Solutions Canada Ltd.