



NEW

# BRAZEN ALL IN®

Post-emergence herbicide for the control of grassy weeds with great tank-mix flexibility and a built-in adjuvant.



## Active Ingredient:

Pinoxaden 50 g/L = EC

## Application Rates & Acres Treated:

Rate: 500 ml/ac

Acres Treated: 20 ac/jug

## Packaging:

Case: 2 x 10 L jugs

## Water Volume:

Ground: 20–40 L/ac  
(5–10 US gal/ac)

Aerial: 12 L/ac  
(3 US gal/ac)

## Rainfastness:

1 hour

## GROUP 1

### REGISTERED CROPS:

- Barley
- Wheat (spring, winter)

### WEEDS CONTROLLED:

- Barnyard Grass
- Green & Yellow Foxtail
- Proso Millet
- Volunteer Canary Seed
- Volunteer Oats
- Wild Oats

### HOW IT WORKS:

BRAZEN ALL IN® Herbicide is absorbed by the leaves and is rapidly translocated to the growing points of leaves and stems. Thorough coverage of the plants is essential for consistent control. Actively growing susceptible grasses stop growing within 48 hours of treatment. Depending on species, growing conditions and crop competition, leaves and growing points turn yellow within one to three weeks after application. Further colour changes and loss of vigour will be observed, followed by a browning and control three to five weeks after application.

### CROP/WEED STAGING:

Crops/Weeds	Growth Stage
Wheat (spring, winter), Barley	1 leaf to flag leaf stage
Wild Oats, Green and Yellow Foxtail, Barnyard Grass, Volunteer Oats, Volunteer Canary Seed, Proso Millet	1–6 leaf, prior to 4 <sup>th</sup> tiller

\* Do not apply past flag leaf stage

### CROP ROTATIONS:

No restrictions.

### GRAZING RESTRICTIONS:

7 days

### PRE-HARVEST INTERVALS:

Grain, Straw: 60 days  
Hay: 30 days

### STORAGE:

If frozen, allow to thaw and agitate thoroughly prior to use.



# BRAZEN ALL IN<sup>®</sup>

## REGISTERED AND SUPPORTED TANK MIXES<sup>1</sup>

### Herbicides:

- BADGE<sup>®</sup>
- Barricade<sup>®</sup> II
- BROMOTRIL<sup>®</sup>
- Cirpreme<sup>™</sup> XC
- Curtail<sup>®</sup> M<sup>2</sup>
- ESTEEM<sup>®</sup>
- ESTEEM ALL IN<sup>®</sup>
- Exhilarate<sup>™</sup>
- FORCEFIGHTER ALL IN<sup>®</sup>
- FORCEFIGHTER M<sup>®</sup>
- Infinity<sup>®</sup>0<sup>2</sup>
- Infinity<sup>®</sup> FX
- MCPA Ester 600<sup>2</sup>
- MCPA Amine (assume 500 series)
- OUTSHINE<sup>®</sup>
- OUTSHINE ALL IN<sup>®</sup>
- Pixxaro<sup>™</sup>
- Pulsar<sup>®</sup>
- Refine<sup>®</sup> SG
- TOPLINE<sup>®</sup>
- Travallas<sup>®</sup>
- Trophy<sup>®</sup>0<sup>2</sup>

### Fungicides:

- BUMPER<sup>®</sup> 432 EC
- TOPNOTCH<sup>™</sup>

<sup>1</sup>Always consult the label of the broadleaf herbicide prior to use.

<sup>2</sup>For control of common ragweed and suppression of round-leaved mallow.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

## MIXING INSTRUCTIONS:

1. Ensure that the sprayer interior is clean, then fill the spray tank with ½ the required amount of water and engage gentle agitation. Good agitation is indicated by a rippling or rolling action on the surface of the water.
2. Add any WG or DF formulation mix partners and agitate to ensure complete mixing.
3. Add any SC formulation mix partners and agitate to ensure complete mixing.
4. Add BRAZEN ALL IN<sup>®</sup> (EC) herbicide and agitate to ensure complete mixing.
5. Add any/additional EC formulation mix partners and agitate to ensure complete mixing.
6. Fill the tank to ¾ the required amount of water.
7. Add any solution (SN) formulation mix partners and agitate to ensure complete mixing.
8. Finish filling the sprayer with water, maintaining good agitation.
9. After any break in spraying operations, agitate thoroughly before spraying again.
10. Spray the herbicide suspension the same day as mixing.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



Always read and follow label directions.  
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

BADGE<sup>®</sup>, BROMOTRIL<sup>®</sup>, BUMPER<sup>®</sup>, ESTEEM<sup>®</sup>, ESTEEM ALL IN<sup>®</sup>, FORCEFIGHTER<sup>®</sup> M, FORCEFIGHTER ALL IN<sup>®</sup>, OUTSHINE<sup>®</sup>, OUTSHINE ALL IN<sup>®</sup>, and TOPLINE<sup>®</sup> are registered trademarks, and TOPNOTCH<sup>™</sup> is a trademark of an ADAMA Group Company © 2025 ADAMA Agricultural Solutions Canada Ltd.

## Quick Tips

Under ideal growing conditions expect to see onset of symptomology in 3-5 days.