



ADAMA

# BRAZEN® II

One of the most trusted and widely used graminicides for grassy weed control in spring wheat and barley with great tank-mix flexibility.



## GROUP 1

### Active Ingredient:

Pinoxaden 100 g/L  
(contains Cloquintocet-mexyl  
at 25 g/L as a safener)

### Application Rates & Acres Treated:

Rate: 160–240 ml/ac

Acres Treated: 40–60 ac/case;  
320–485 ac/drum

### Packaging:

Case: BRAZEN® II: 1 × 9.7 L jug;  
Cohere™ Adjuvant: 1 × 11.3 L jug

Bulk: BRAZEN® II: 77.6 L drums;  
Cohere™ Adjuvant: 90.4 L drum

### Water Volume:

Ground: 20–40 L/ac (5–10 US gal/ac)  
Aerial: 12 L/ac (3 US gal/ac)

### Rainfastness:

1 hour

### MIXING INSTRUCTIONS:

1. Clean spray tank and ½ fill with clean water. Start agitation or bypass system.
2. If a broadleaf herbicide is to be used, add the product first prior to adding BRAZEN® II and agitate for 2–3 minutes.
3. Add correct amount of BRAZEN® II.
4. Agitate for 2–3 minutes.
5. Add correct amount of Cohere™ Adjuvant.
6. Agitate for 1–2 minutes before adding remainder of water and then maintain constant agitation.
7. After any break in spraying operations, agitate thoroughly before spraying again.
8. Use the spray suspension as soon as it is prepared.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

### HOW IT WORKS:

BRAZEN® II is absorbed by the leaves and is rapidly translocated to the growing points of leaves and stems. Thorough coverage of the plants is essential for consistent control. Actively growing susceptible grasses stop growing within 48 hours of treatment. Depending on species, growing conditions and crop competition, leaves and growing points turn yellow within 1–3 weeks after application. Further colour changes and loss of vigour will be observed, followed by a browning and control 3–5 weeks after application.



# BRAZEN® II

## CROP STAGING:

Crops/Weeds	Growth Stage
Barley, Spring Wheat	1 leaf to flag leaf stage
Green And Yellow Foxtail, Persian Darnel, Proso Millet, Volunteer Canary Seed, Volunteer Oats, Wild Oats	1 – 6 leaf, prior to 4 <sup>th</sup> tiller

(BRAZEN® II is not registered for use on durum wheat)

## REGISTERED AND SUPPORTED TANK MIXES<sup>1</sup>

### Herbicides:

- BADGE®
- Barricade® II
- BROMOTRIL®
- Cirpreme™ XC
- Curtail® M<sup>4</sup>
- ESTEEM®
- ESTEEM ALL IN®
- Exhilarate™
- FORCEFIGHTER ALL IN®
- FORCEFIGHTER® M
- Infinity®<sup>4</sup>
- Infinity FX®
- MCPA Ester 600<sup>4</sup>
- MCPA Amine
- OUTSHINE®
- OUTSHINE ALL IN®
- Pixxaro®
- Pulsar®
- Refine® SG
- TOPLINE®
- Travallas®
- TROPHY®<sup>4</sup>

### Fungicides:

- BUMPER® 432 EC
- TOPNOTCH™

<sup>1</sup>Always consult the label of the broadleaf herbicide prior to use.

<sup>2</sup>Addition of surfactants other than ADAMA Adjuvant 80 are not required.

<sup>3</sup>Suppression only of green foxtail.

<sup>4</sup>For control of common ragweed and suppression of round-leaved mallow.

## ADJUVANT RATE

283 ml/ac; adjuvant co-pack

## GRAZING RESTRICTIONS:

7 days

## CROP ROTATIONS:

No restrictions.

## STORAGE:

If frozen, allow product to thaw and agitate thoroughly prior to use.

## PRE-HARVEST INTERVALS:

Grain, Straw: 60 days

Hay: 30 days

[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



Always read and follow label directions.  
Toll-free: 1.855.264.6262  
Website: ADAMA.COM

BADGE®, BRAZEN®, ESTEEM®, ESTEEM ALL IN®, FORCEFIGHTER ALL IN®, FORCEFIGHTER® M, INVOLVE® 50 WDG, OUTSHINE®, OUTSHINE ALL IN®, and TOPLINE® are registered trademarks, and TOPNOTCH® is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.