



ADAMA

HERBICIDE



NEW

GORDEX™

Power up your pre-seed cereal burnoff: Tank-mix the multi-mode liquid formulation of GORDEX™ with glyphosate for extended residual control of Volunteer Canola and effective knockdown of tough Group 2 & 9 resistant weeds—like Kochia.

GROUP 2 & 4

Active Ingredient:

Florasulam 200 g/L = SC
Dicamba 480 g/L (DGA salt) = SL

Application Rates:

SPRING or FALL RATE: GORDEX™ 160 ac/case

- GORDEX™ A @ 0.007 L/ac
(160 ac/1.13 L jug)
- GORDEX™ B @ 0.097 L/ac
(80 ac/7.76 L jug)

FALL RATE only: GORDEX™ 112 ac/case

- GORDEX™ A @ 0.01 L/ac
(112 ac/1.13 L jug)
- GORDEX™ B @ 0.138 L/ac
(56 ac/7.76 L jug)

Packaging

& Acres Treated:

160 ac/co-pack (spring or fall rate);
112 ac/co-pack (fall rate only)

GORDEX™ A = 1.13 L (florasulam)

GORDEX™ B = 2 x 7.76 L
Dicamba 480 SC (DGA salt)

Acres Treated: 112 – 160 ac/co-pack case

Water Volume:

Ground: 20 – 40 L/ac (5 – 10 US gal/ac)
Aerial: Do not apply

Rainfastness:

30 minutes

KEY BENEFITS:

- First co-pack offering with two liquid formulations and convenient packaging
- Multi-mode of action tank mix partner with glyphosate for pre-seed cereals
- Offers extended control of Volunteer Canola + contact control of group 2/9 resistant Kochia

REGISTERED CROPS:

For a complete list of all crops registered for GORDEX™ please refer to the product labels

- Oats
- Spring Barley
- Spring Wheat

WEEDS CONTROLLED:

Weeds controlled when tank mixed with glyphosate (0.5 REL/ac)

Controlled (2-4 leaf stage):

- | | | |
|--------------------------------|---|---------------------------------|
| • Canada Fleabane ¹ | • Hemp-Nettle | • Shepherd's Purse |
| • Cleavers | • Kochia | • Smartweed |
| • Common Chickweed | • Lady's Thumb | • Stinkweed |
| • Common Ragweed ¹ | • Lamb's Quarters | • Volunteer Canola ³ |
| • Cow Cockle | • Narrow-Leaved Hawk's Beard ¹ | • Volunteer Flax |
| • Dandelion ² | • Redroot Pigweed | • Volunteer Wheat |
| • Flixweed | • Russian Thistle | • Wild Buckwheat ⁴ |
| | | • Wild Mustard |

Grass weeds controlled:

- | | | |
|-----------------|------------------|--------------------|
| • Downy Brome | • Green Foxtail | • Volunteer Barley |
| • Giant Foxtail | • Persian Darnel | • Wild Oats |

ADDITIONAL WEEDS CONTROLLED WHEN TANK MIXED WITH GLYPHOSATE @ 1-2.8 REL/AC

Controlled (2-4 leaf stage):

- | | |
|-------------------------------|--------------------------------------|
| • Annual Sow Thistle | • Perennial Sow Thistle ⁵ |
| • Canada Thistle ⁶ | • Quackgrass |

¹ Less than 3 inches in height

² Mature plants up to 12 inches in diameter, rosettes, and seedlings

³ Including all herbicide-tolerant canola varieties

⁴ Up to 5 leaves

⁵ Advanced staged applications will be less effective

⁶ Rosette Stage

GORDEX™


HOW IT WORKS:

Provides extended pre-seed control of several broadleaf weeds which helps reduce flushing of weeds and protects critical weed free period of your cereal crop.

APPLICATION TIMING & RE-CROPPING:

Spring: Apply prior to seeding. Do not apply later than 48 hours post seeding.

Fall: Post harvest – prior to freeze up.



MIXING INSTRUCTIONS:

1. Fill the clean spray tank ½ full of clean water. Start agitation system.
2. Add the required amount GORDEX™ A. Continue to agitate.
3. Add the required amount of GORDEX™ B. Continue to agitate.
4. Add required amount of glyphosate.*
5. Complete filling with remaining amount of water and continue agitation.

*Can be tank mixed with all forms of glyphosate. Do not mix different glyphosate salts (DMA, IPA or k+) together.

REGISTERED & SUPPORTED TANK MIXES:

- Glyphosate

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions for Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

CROP ROTATIONS:

When applied prior to August 1: Barley, Canola, Oats, Field Peas, and Wheat (Spring & Durum) can be seeded the following year.

When applied after August 1: Only Wheat (Spring & Durum), Barley, and Oats can be seeded the following year.

PRE-HARVEST INTERVALS:

Do not harvest the treated crop within 60 days of application.

GRAZING RESTRICTIONS:

- Livestock may graze fields within 7 days after application.
- DO NOT harvest forage or cut within 60 days after application.

When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions and grazing limitations for each product used in the tank-mix.

STORAGE:

- Store GORDEX™ in its original container only, away from other pesticides, fertilizer, food, or feed.
- Keep the container closed to prevent spills and contamination.
- Keep packages dry at all times.

[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



QUICK TIPS:

Use multi-mode of action GORDEX™ as a glyphosate tank-mix before key cereals in spring or fall for superior control of problem weeds such as Kochia, Wild Buckwheat, and Dandelion, plus extended control of Volunteer Canola and other weeds.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

GORDEX™ is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2025 ADAMA Agricultural Solutions Canada Ltd.