



ADAMA

EMPHASIS® MAX

EMPHASIS® MAX with its dual mode of action, has an increased rate of carfentrazone for enhanced control of Kochia, including Group 2, 4, 5 and 9 resistant biotypes.



Active Ingredients:

EMPHASIS® A : Carfentrazone-ethyl at 240 g/L = EC, BROMOTRIL® 240 EC: Bromoxynil Octanoate Ester at 240 g/L = EC

Application Rates:

Canola:

Rate: 30 ml/ac EMPHASIS® A
+ 236 ml/ac BROMOTRIL® 240 EC

Wheat, Barley and Oats:

Standard Rate: 30 ml/ac of EMPHASIS® A
+ 472 ml/ac of
BROMOTRIL® 240 EC

Packaging and Acres Treated:

Co-pack: 2 × 1.2 L EMPHASIS® A
+ 2 × 9.7 L BROMOTRIL® 240 EC

Canola Acres Treated: 80 ac/case, can be tank mixed with glyphosate

Wheat, Barley, Oats Acres Treated: 40 ac/case can be tank mixed with glyphosate, with a 1.2 L jug EMPHASIS® A left over

Water Volume:

Ground: 40 L/ac (10 US gal/ac)

Aerial: Do not apply by air.

Rainfastness:

6 hours

GROUP 6 & 14

REGISTERED CROPS AND STAGING:

Crop	Stage
Barley, Oats, Wheat, Canola	Pre-seed burn-off with glyphosate

HOW IT WORKS:

EMPHASIS® MAX is a multi mode contact herbicide that controls broadleaf weeds including Group 2 and 9 resistant biotypes such as Kochia. Within a few hours following application, the foliage of susceptible weeds shows signs of desiccation, and in subsequent days, necrosis and death of the plant occur.

MIXING INSTRUCTIONS:

1. Fill spray tank with ½ of the volume of clean water needed.
2. With agitator running add the required amount of EMPHASIS® A to spray tank.
3. Next add the required amount of BROMOTRIL® 240 EC.
4. Add more water, then add glyphosate.
5. Complete filling the tank to desired level.

ADJUVANT RATE:

No adjuvant required.

CROP ROTATIONS:

No restrictions.

STORAGE:

- Do not freeze.



EMPHASIS® MAX

WEEDS CONTROLLED:

When used as directed, EMPHASIS® A will provide control of the listed weeds up to ten (10) cm in height, or as specified. See the individual labels of each tank-mix partner for specific rates and weed staging and always follow the directions for use of the most stringent label. Good spray coverage is essential for optimal weed control.

Weeds Controlled	Rates by Product	
	EMPHASIS® A Rate	BROMOTRIL® 240 EC Rate
EMPHASIS® MAX alone:	80 ac/case - Pre-Seed Canola	
Black Nightshade (up to 5 cm), Kochia, Lamb's Quarters (up to 7.5 cm), Redroot Pigweed, Round-Leaved Mallow, Russian Thistle (up to 5cm), Shepherd's Purse, Stinkweed, Tall Waterhemp (up to 5 cm), Volunteer Canola	30 ml/ac	236 ml/ac
American Nightshade, Bluebur, Carpetweed, Common Buckwheat, Common Groundsel, Common Purslane, Common Waterhemp, Hairy Nightshade, Jimsonweed, Pale Smartweed, Tansy Mustard, Tartary Buckwheat, Tumble Pigweed	40 ac/case - Pre-Seed Wheat, Barley & Oats	
	30 ml/ac 1 jug of EMPHASIS® A will remain	472 ml/ac
EMPHASIS® MAX + Glyphosate	Glyphosate REL/ac	Glyphosate Grams of a.i./ac
Weeds listed above by rate + Cocklebur, Cow Cockle, Green Foxtail, Green Smartweed, Lady's Thumb, Smooth Pigweed, Volunteer Barley, Volunteer Wheat, Wild Mustard, Wild Oats	0.5 REL/ac	180g a.i./ac
Weeds Listed Above + Common Ragweed, Wild Buckwheat, Canada Fleabane, Cleavers, Downy Brome, Flixweed, Giant Foxtail, Hemp-Nettle, Persian Darnel, Russian Thistle, Stinkweed, Volunteer Flax, Narrow-Leaved Hawk's Beard	0.75 REL/ac	277g a.i./ac
Weeds Listed Above + Annual Bluegrass, Annual Sow Thistle, Canada Thistle (rosette stage, summerfallow), Crabgrass, Dandelion (less than 15 cm), Kochia, Narrow-Leaved Vetch, Prickly Lettuce, Quackgrass (light to moderate infestations, 3–4 green leaves or more), Shepherd's Purse	1 REL/ac	360g a.i./ac

*Refer to the glyphosate label for the complete list of weeds controlled at each rate; the EMPHASIS® MAX + glyphosate combination controls ~70 weeds, not all listed here.



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

BROMOTRIL® and EMPHASIS® MAX are registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.

Quick Tips

Wait at least 1 day after application before seeding. Allow adequate time for weed control. Avoid overnight storage of spray mixtures when possible. Check weather conditions in advance. Premixing EMPHASIS® MAX spray solutions in nurse tanks is not recommended. Use a water volume of at least 40 L/ac (10 US gal/ac) for adequate coverage.