



BRAZEN® II

One of the most trusted and widely used
graminicides for grassy weed control in spring
wheat and barley with great tank-mix flexibility.



GROUP 1

Active Ingredient:

Pinoxaden 100 g/L

Application Rates:

Rate: 160–240 ml/ac

Packaging and

Acres Treated:

Case: BRAZEN® II: 1 x 9.7 L jug;
Cohere™ Adjuvant: 1 x 11.3 L jug

Drum: BRAZEN® II: 77.6 L drums;
Cohere™ Adjuvant: 90.4 L drum

Acres Treated: 40–60 ac/case;
320–485 ac/drum

Water Volume:

Ground: 20–40 L/ac (5–10 US gal/ac)

Aerial: 12 L/ac (3 US gal/ac)

Rainfastness:

1 hour



FOR FULL PRODUCT
DETAILS, SCAN THE
QR CODE.

REGISTERED CROPS:

- Barley
- Spring Wheat

WEEDS CONTROLLED:

| | | |
|------------------|-------------------------|------------------|
| • Barnyard Grass | • Proso Millet | • Volunteer Oats |
| • Green Foxtail | • Volunteer Canary Seed | • Wild Oats |
| • Persian Darnel | | • Yellow Foxtail |

CROP STAGING:

| Crops/Weeds | Growth Stage |
|--|---|
| Barley, Spring Wheat | 1 leaf to flag leaf stage |
| Barnyard Grass, Green And Yellow Foxtail, Persian Darnel, Proso Millet, Volunteer Canary Seed, Volunteer Oats, Wild Oats | 1–6 leaf, prior to 4 th tiller |

HOW IT WORKS:

BRAZEN® II is absorbed by the leaves and is rapidly translocated to the growing points of leaves and stems. Thorough coverage of the plants is essential for consistent control. Actively growing susceptible grasses stop growing within 48 hours of treatment. Depending on species, growing conditions and crop competition, leaves and growing points turn yellow within 1–3 weeks after application. Further colour changes and loss of vigour will be observed, followed by a browning and control 3–5 weeks after application.

BRAZEN® II

REGISTERED AND SUPPORTED TANK MIXES¹:

Herbicides:

- BADGE®
- Barricade® II
- BROMOTRIL® 240 EC
- Cirpreme™ XC
- Curtail® M²
- ESTEEM ALL IN®
- Exhilarate™
- FORCEFIGHTER ALL IN®
- Infinity®²

- Infinity FX®
- MCPA ESTER 600²
- MCPA Amine (assume 500 series)
- OUTSHINE ALL IN®
- Pixxaro®
- Pulsar®
- Refine® SG
- Travallas®
- Trophy®²

Fungicides:

- BUMPER® 432 EC
- TOPNOTCH™

¹Always consult the label of the broadleaf herbicide prior to use.

²For control of Common Ragweed and suppression of Round-Leaved Mallow.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

1. Clean spray tank and ½ fill with clean water. Start agitation or bypass system.
2. If a broadleaf herbicide is to be used, add the product first prior to adding BRAZEN® II and agitate for 2–3 minutes.
3. Add correct amount of BRAZEN® II.
4. Agitate for 2–3 minutes.
5. Add correct amount of Cohere™ Adjuvant.
6. Agitate for 1–2 minutes before adding remainder of water and then maintain constant agitation.
7. After any break in spraying operations, agitate thoroughly before spraying again.
8. Use the spray suspension as soon as it is prepared.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

ADJUVANT RATE:

283 ml/ac; adjuvant co-pack

GRAZING RESTRICTIONS:

7 days

CROP ROTATIONS:

No restrictions.

STORAGE:

If frozen, allow product to thaw and agitate thoroughly prior to use.

PRE-HARVEST INTERVALS:

Grain, Straw: 60 days

Hay: 30 days



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

BADGE®, BRAZEN®, BROMOTRIL®, BUMPER®, ESTEEM ALL IN®, FORCEFIGHTER ALL IN® and OUTSHINE ALL IN® are registered trademarks, and TOPNOTCH™ is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

Apply to actively growing weeds for best results. An early application will maximize crop yields by reducing weed competition. Weeds emerging after application of BRAZEN® II will not be controlled.