



2,4-D ESTER 700

Provides reliable control of broadleaf weeds and great tank-mix flexibility in wheat, barley, rye and other crops.



GROUP 4

Active Ingredient:

2,4-D Ester 2 EH Ester
660 g/L = EC

Application Rates:

Rate: 200 – 600 ml/ac

Packaging and Acres Treated:

Case: 2 x 10 L jugs
Bulk: 120 L drums
Tote: 500 L
MAX Tote: 1000 L

Acres Treated: 17 – 50 ac/jug;
200 – 600 ac/drum;
850 – 2,500 ac/tote;
1,665 – 5,000 ac/tote

Water Volume:

Ground: 20 – 60 L/ac
(5 – 15 US gal/ac)

Aerial: Minimum 12 L/ac (3 US gal/ac)

Rainfastness:

2 hours

REGISTERED CROPS:

- Barley
- Field Corn
- Rye (spring, fall)
- Wheat (spring, winter)

See registered label for timing and rates for each crop.

WEEDS CONTROLLED:

Susceptible weeds:

- Annual Sow Thistle¹
- Bluebur
- Burdock¹
- Cocklebur
- Daisy Fleabane
- False Flax
- False Ragweed
- Flixweed
- Giant Ragweed
- Goat's Beard
- Kochia
- Lamb's Quarters
- Mustards (except dog and tansy)
- Narrow-Leaved Hawk's Beard²
- Plantain
- Prickly Lettuce
- Ragweed
- Redroot Pigweed
- Russian Pigweed
- Russian Thistle
- Shepard's Purse
- Stinging Nettle
- Stinkweed
- Sweet Clover (seedling)
- Thyme-Leaved Spurge
- Volunteer Canola¹
- Wild Radish
- Wild (prairie) Sunflower

Harder-to-control weeds:

- Curled Dock¹
- Dog Mustard
- Field Peppergrass
- Flixweed⁴
- Groundsel
- Hairy Galinsoga
- Hawkweed
- Heal-All
- Knotweed¹
- Narrow-Leaved Hawk's Beard⁴
- Oak-Leaved Goosefoot
- Pineapple Weed
- Prostrate Pigweed
- Purslane
- Sheep Sorrel
- Tansy Mustard
- Tumble Pigweed
- Velvetleaf
- Volunteer Canola^{1,3}

¹All types

²In the fall, and at the 1- to 2-leaf stage in spring

³4- to 6-leaf stage

⁴If treated before bolting in spring



2,4-D ESTER 700

WEEDS CONTROLLED:

Very-hard-to-control weeds:

- Biennial Wormwood
- Blue Lettuce
- Bull Thistle
- Burdock
- Buttercup
- Canada Thistle
- Chicory
- Curled Dock
- Dandelion
- Field Bindweed
- Field Chickweed²
- Field Horsetail²
- Gumweed
- Hedge Bindweed
- Hemp-Nettle²
(if treated before the 4-leaf stage)
- Hoary Cress
- Lady's Thumb²
- Leafy Spurge
- Mouse-Eared Chickweed²
- Perennial Sow Thistle
- Russian Knapweed
- Scentless Mayweed
- Smartweed²
- Tartary Buckwheat
- Teasel
- Volunteer Sunflower
- Wild Buckwheat²
- Yellow Rocket¹

¹ 1- to 3-leaf stage

² Use highest listed rate for suppression

HOW IT WORKS:

Systemic activity hinders plant cell growth in newly forming stems and leaves promoting uncontrolled, unsustainable growth, causing stem curl-over, leaf withering and eventual plant death.

REGISTERED AND SUPPORTED TANK MIXES:

- BISON® 400 L
- BUMPER® 432 EC
- PYRINEX® 480 EC
- Traxos®

PRE-SEED:

- BROMOTRIL® 240 EC
- Glyphosate
- INVOLVE® 50 WDG

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

1. ½ fill the tank with clean water.
2. Add the required amount of ADAMA 2,4-D Ester 700 and agitate thoroughly.
3. Add any tank-mix partners.
4. Fill the tank and agitate again before use.

PRE-HARVEST INTERVAL:

90 days

CROP ROTATIONS:

No restrictions.

GRAZING RESTRICTIONS:

30 days

STORAGE:

May be stored at any temperature.
Shake well before using.



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

BISON®, BROMOTRIL®, BUMPER®, INVOLVE® and PYRINEX® are registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

Avoid spray drift to any off-target vegetation.
Coarse sprays are less likely to drift. Do not spray during periods of high winds.