



ADAMA

BADGE®

Multi-mode of action tank-mix partner
for cereals, flax and corn.

GROUP 4 & 6**Active Ingredients:**

Bromoxynil 225 g/L and
MCPA 225 g/L = EC

Application Rates:

Rate: 500 ml/ac

**Packaging and
Acres Treated:**

Case: 2 x 10 L jugs

Bulk: 120 L

Tote: 450 L

Acres Treated: 20 ac/jug;
240 ac/drum;
900 ac/tote

Water Volume:**Ground:**

Cereals and flax: 20 – 40 L/ac
(5 – 10 US gal/ac);

Corn: 80 – 120 L/ac
(20 – 30 US gal/ac);

Forages: 60 L/ac
(15 US gal/ac)

Aerial (wheat, barley and oats only):
8 – 20 L/ac (2 – 5 US gal/ac)

Rainfastness:

1 hour

REGISTERED CROPS:

- Barley
- Canary Seed
- Corn
- Fall Rye
- Flax (including low linolenic varieties)
- Oats
- Seedling Grasses
- Timothy (established for seed production)
- Wheat (spring, winter, durum)

WEEDS CONTROLLED:

- American Nightshade
- Ball Mustard
- Bluebur
- Canada Thistle¹
- Cocklebur
- Common Buckwheat
- Common Groundsel
- Common Ragweed
- Cow Cockle²
- Flixweed
- Green Smartweed
- Kochia³
- Lady's Thumb
- Lamb's Quarters
- Night-Flowering Catchfly
- Pale Smartweed
- Perennial Sow Thistle¹
- Redroot Pigweed
- Russian Thistle³
- Scentless Chamomile⁴
- Shepherd's Purse
- Stinkweed
- Tartary Buckwheat
- Velvetleaf⁵
- Volunteer Canola (all types)
- Volunteer Sunflower
- Wild Buckwheat
- Wild Mustard
- Wild Tomato
- Wormseed Mustard

¹Top growth control.

²Up to 4-leaf stage.

³Spray before plants are 2 inches high.

⁴Spring annual only.

⁵Spray before plants are 3 inches high.

BADGE®

HOW IT WORKS:

A combination of systemic and contact activity with weeds yellowing within 2–4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2–10 days.

CROP STAGING:

Crop	Stage
Barley, Oats, Wheat (spring, durum)	2 leaf to early flag
Canary Seed	3–5 leaf
Corn	4–6 leaf
Fall Rye	When growth commences in spring to early flag leaf.
Flax (including low linolenic varieties)	2 inches to early bud stage. Best tolerance occurs when flax is 2–4 inches tall.
Seedling Grasses	2–4 leaf
Timothy (established for seed production)	Prior to shot blade in the seed production year.
Winter Wheat	2–4 leaf stage in the fall or after growth resumes up to early flag leaf.

REGISTERED AND SUPPORTED TANK MIXES:

- Corn: Atrazine
- Flax: ARROW ALL IN® or Poast®
- Oats: MCPA Ester 600
- Spring Wheat and Barley: Ally®, MCPA Ester 600, Refine® SG, BISON® 400 L
- Spring Wheat only: LADDER ALL IN®, Everest® 3.0 or Traxos®
- Winter Wheat: MCPA Ester 600, Everest® 3.0 or Refine® SG

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

MIXING INSTRUCTIONS:

1. Fill clean spray tank ½ full with water.
2. Add the required amount of BADGE® and agitate thoroughly.
3. Fill the tank and agitate again before use.
4. If tank mixing with a grassy weed herbicide, read both labels and follow the more stringent directions for tank mixing.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

CROP ROTATIONS:

No re-cropping restrictions the year after treatment.

PRE-HARVEST INTERVAL:

Flax: 60 days

STORAGE:

Do not freeze.

GRAZING RESTRICTIONS:

Do not graze treated grain or established Timothy crops or cut for feed within 30 days of application.



[CLICK HERE FOR FULL PRODUCT DETAILS.](#)



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

ARROW®, ARROW ALL IN®, BADGE®, BISON® and LADDER ALL IN® are registered trademarks of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

BADGE® herbicide is well known for being gentle on the crop. Avoid spraying if temperatures are above 25°C.