



ADAMA

NEW

OXALIS®

An effective resistance management tank mix partner, OXALIS® is a pre-seed soil applied herbicide that gives extended control of key broadleaf weeds prior to their emergence.

**Active Ingredient:**

Flumioxazin 480 g/L = SC

Application Rates:

Rate: 60–90 ml/ac

Packaging & Acres Treated:

Case: 2 x 4.8 L

Acres Treated: 106–160 acres/case

Water Volume:

Ground: 40–120 L/ac

(10–30 US gal/ac)

Aerial: Do not apply

GROUP 14**REGISTERED CROPS:**

- Field Peas
- Soybeans
- Field Corn

KEY BENEFITS:

- Residual control of later-emerging weed flushes and extend protection for up to 8 weeks after application.
- Provides a clean start by controlling susceptible broadleaf weeds during the early stages of crop development.
- Strengthen resistance management programs by introducing a Group 14 mode of action to control key broadleaf weeds, including Kochia resistant to Groups 2, 4, and 9.

WEEDS CONTROLLED:

- Canada Fleabane
- Common Ragweed
- Common Lamb's-Quarters
- Dandelion
- Eastern Black Nightshade
- Green Foxtail¹
- Green Pigweed
- Hairy Nightshade
- Kochia
- Redroot Pigweed

¹ Suppression only.**WEED STAGING :**

- When weeds are already emerged, apply OXALIS® as a tank mix with a glyphosate product at 364–485 g a.e./ac.

CROP STAGING:

- Field Peas: (for Fall burndown) – Apply in the fall, just before freeze-up and when winter annuals and perennial weeds are still growing.
- Field Corn: (Spring and Fall application) – Apply 7 to 30 days prior to planting field corn into no-till or minimum tillage fields.
- Soybeans*: (Spring and Fall application) – Apply prior to planting or within 3 days after planting and prior to soybean emergence.

*No-till planter that incorporates the soil during planting may result in decreased weed control in the row—recommendation is to apply OXALIS® within 3 days of planting before soybeans emerge.

CROP	SOIL TYPE ¹	RATE
Field Corn, Field Peas, Soybeans	Coarse- textured, with <5% organic matter	60 ml/ac
	Medium- and fine-textured, with <5% organic matter	90 ml/ac

¹ Do not apply on soils with > 5% organic matter.

**CLICK HERE FOR
FULL PRODUCT
DETAILS.**

OXALIS®

HOW IT WORKS:

OXALIS® controls weeds by inhibiting protoporphyrinogen oxidase (PPO), an essential enzyme required for normal plant growth. Susceptible weed seedlings absorb the herbicide during emergence, and exposure to sunlight triggers rapid cell membrane damage, resulting in weed death.

MIXING INSTRUCTIONS:

1. Fill clean spray tank ½ to ¾ of desired level with clean water. Engage gentle agitation.
2. While agitating, slowly add OXALIS® to the spray tank. Agitation should create a rippling or rolling action on the water surface. Add tank mix partner if required following WAMLEGS.
3. Add adjuvants or surfactants, if recommended.
4. Fill spray tank to desired level with water. Agitation should continue until spray solution has been applied.
5. Mix only the amount of spray solution that can be applied the day of mixing. OXALIS® should be applied within 6 hours of mixing.

ROTATIONAL RESTRICTIONS:

The following rotational crops may be planted after applying OXALIS® at the listed rate. Planting earlier than the recommended rotational interval may result in crop injury.

RATE	CROP	ROTATIONAL INTERVAL
60 ml/ac	Soybean	Immediately
	Field Corn	7 days
	Sorghum, Dry Common Beans ¹ , and Canola	9 months
	Alfalfa and Barley	11 months
	All other crops not listed ²	12 months
90 ml/ac	Soybean	Immediately
	Field Corn (minimum and no-till)	14 days
	Field Corn (conventional till)	30 days
	Sorghum, Dry Common Beans ¹	9 months
	Alfalfa, Barley, and Canola	11 months
	All other crops not listed ²	12 months

¹ Common bean tolerance to OXALIS® varies by variety. Test a small area first and consult your seed supplier before broad adoption.

² Successful soil bioassay must be performed prior to planting crops not listed.

After periods of extended drought, when there is a lack of adequate soil moisture, longer rotational intervals may be needed. A successful soil bioassay should be performed prior to planting potential rotational crops to confirm safety.

GRAZING RESTRICTION:

DO NOT graze, cut or feed treated crops to livestock.

REGISTERED AND SUPPORTED TANK MIXES:

- Glyphosate

ADJUVANT:

Not all tank mixes require an adjuvant, see label for details.

STORAGE:

Store in a cool, dry, secure place.

QUICK TIPS:

Fall applications should be made just before freeze-up. Do not apply to snow covered or frozen soil.

Do not incorporate or perform any tillage operations after application or weed control will be reduced.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

OXALIS® is a registered trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.