

**NEW****SAFACYN®****Pre-Harvest/Harvest Aid**

A new standard in harvest aid and weed management. This advanced formulation delivers fast and complete crop and weed dry down, while also enabling flexibility to use the adjuvant type and rate that fits your field & agronomic needs.

HERBICIDE**Active Ingredient:**

Saflufenacil 160 g/L SL

Application Rates:

Rate: 63–92 ml/ac of
SAFACYN® + adjuvant

Packaging**& Acres Treated:**

Case: 2 x 7.35 L

Acres Treated: 160–233 acres

Water Volume:

Ground: 40–80 L/ac
(10–20 US gal/ac)

Aerial: 20 L/ac (5 US gal/ac)

Rainfastness:

Avoid application when heavy
rain is in the forecast.



**CLICK HERE
FOR FULL
PRODUCT
DETAILS.**

GROUP 14**KEY BENEFITS:**

- **Faster, more complete crop and weed dry down:** Delivers consistent dry down for a faster, more efficient harvest.
- **Flexible adjuvant options:** Choose the adjuvant type and rate that best fits your crop, weed pressure and environmental conditions.
- **Improved harvest efficiency:** Supports easier cutting, faster combining, reduced fuel use, and lower dockage.
- **Flexible fit across multiple cropping systems:** Fits with all canola herbicide systems and is registered across a wide range of crops including canola, cereals, peas, lentils, soybeans, and flax.
- **Cleaner fields next spring:** Helps manage late-season weeds and sets fields up for a cleaner start next season.

HARVEST AID & PRE-HARVEST WEED MANAGEMENT:

SAFACYN® may be used as a harvest aid to accelerate the rate of crop dry down and improve crop uniformity to facilitate direct combining. Early application may result in yield and/or seed quality loss.

SAFACYN® may be used as a pre-harvest weed management in wheat (including durum, spring and winter wheat), barley (including feed varieties), triticale to improve dry down of Volunteer Canola (all types including Roundup Ready®), Common Ragweed, Canada Fleabane, Redroot Pigweed and Wild Buckwheat and to facilitate direct combining. Early application may result in yield loss.

CROP	TIMING	Seed Production or Restrictions on Glyphosate Use	No Restrictions on Glyphosate Use
Canola, Mustard	Apply when 80% of seeds have changed colour.	92 ml/ac of SAFACYN® (160ac/case) + adjuvant	92 ml/ac of SAFACYN® (160 ac/case) + 360 g a.e./ac of glyphosate + adjuvant
Chickpeas	For Desi type, apply at the time swathing would normally begin, when the majority of plants are yellow and most pods are mature and seeds have turned from green to yellow or brown. The upper part of the plant may still be green. For Kabuli type, apply when the majority of plants and pods are ripe and dry with seeds turned from green to white or tan, and detached from the pods. Dry down is less complete in Kabuli type due to its thick pod wall.		
Dry Common Beans, Soybeans	Apply when stems are green to brown in colour and pods are mature (yellow – brown) and 80 – 90% of the original leaves have dropped.		
Faba Beans	Apply when 80% of lower pods have turned black, middle pods have turned yellow/tan, and top green pods have firm seed.		
Flax	Apply when 75% of bolls have turned colour.		
Lentils (red)	Apply when bottom 15% of pods are brown and rattle when shaken.		
Peas	Apply when the majority of pods are brown (70–80%).		
Sunflowers	Apply when the backs of the heads and bracts are turning yellow, and seed moisture is 20–30%.		
Wheat & Barley	Hard dough stage; a thumbnail impression remains on seed; less than 30% moisture.		63–92 ml/ac of SAFACYN® (160–233 ac/case) + 360 g a.e./ac of glyphosate + adjuvant

SAFACYN®

HOW IT WORKS:

SAFACYN® is quickly absorbed through both the roots and leaves of plants and moves throughout the weed once inside. It works by blocking an important enzyme called protoporphyrinogen oxidase (PPO), which is essential for normal plant function. When PPO is inhibited, the weed cell membranes are damaged, causing the plant to die. Under active growing conditions, susceptible weeds can show symptoms within hours of application, with complete plant death typically occurring within 3–5 days depending on environmental conditions. Under cooler or less ideal conditions full kill may take longer.

ADJUVANT RATE:

1. When Mixing SAFACYN® with Glyphosate: use a Methylated seed oil (MSO) or Crop Oil Concentrate (COC) adjuvant at 0.5 – 1% v/v OR a non-ionic surfactant (NIS) at 0.25% v/v.
2. When using SAFACYN® without Glyphosate: use a Methylated seed oil (MSO) or Crop Oil Concentrate (COC) adjuvant at 0.5 – 1% v/v.

REGISTERED AND SUPPORTED TANK MIXES:

- Glyphosate

GRAZING RESTRICTION:

- Canola, Wheat, Barley & Triticale: Straw can be used as feed or grazed 3 days after pre-harvest application.
- Dry Beans, Soybeans: Do not graze or feed treated hay or straw to livestock.

MIXING INSTRUCTIONS:

1. Fill clean tank $\frac{1}{2}$ to $\frac{3}{4}$ full of clean water and turn agitation on.
2. Add the required amount of SAFACYN® herbicide and continue to agitate.
3. If required, add the correct amount of tank-mix partner while agitating.
4. Add the required amount of adjuvant while agitating.
5. Continue agitating and fill the remainder of the spray tank with water.

NOTE: If tank-mixing, please reference the label of the partner for specific mixing order or follow WAMLEGS or WALES for proper mixing protocol.

PRE-HARVEST INTERVAL:

Barley, Canola, Flax, Lentils, Mustard, Peas, Soybeans, Triticale, Wheat: 3 days

Chickpeas, Dry Common Beans, Faba Beans: 2 days

Sunflowers: 7 days

For tank mixtures that include glyphosate please refer to and follow glyphosate pre-harvest interval on label.

STORAGE:

Do not freeze.



Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

SAFACYN® is a registered trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

Water volume is important to ensure good coverage and even dry down of plant tissue. Use the higher end of the recommended water volume range in dense or uneven crops to improve coverage deep into the crop canopy and on late green material.