

HERBICIDE

ADAMA

PRIORITY® HL

The ideal glyphosate tank-mix partner for pre-seed burn-off with proven extended control of non-Clearfield® Volunteer Canola and winter annuals. Upgraded with a convenient high load formulation!

High Load
FORMULATION TECHNOLOGY

GROUP 2

Active Ingredient:

Florasulam 200 g/L = SC

Application Rates:

Rate: 10 ml/ac

Packaging and Acres Treated:

Case: 4 x 1.62 L jugs

Acres Treated: 160ac/jug (640 ac case)

Water Volume:

Ground: 20-40 L/ac (5-10 US gal/ac)

Aerial: Do not apply

Rainfastness:

30 minutes

REGISTERED CROPS:

- Wheat (spring, durum, winter)
- Spring Barley
- Oats

WEEDS CONTROLLED BY PRIORITY® HL + GLYPHOSATE AT 0.5 REL/AC:

Controlled (2–4 leaf stage):

- Canada Fleabane[2]
- Cleavers
- Common Chickweed
- Common Ragweed[2]
- Cow Cockle
- Dandelion[3]
- Flixweed
- Hemp-Nettle
- Lady's Thumb
- Lamb's Quarters
- Narrow-Leaved Hawk's Beard[2]
- Redroot Pigweed
- Russian Thistle
- Shepherd's Purse
- Smartweed
- Stinkweed
- Volunteer Canola[4]
- Volunteer Flax
- Volunteer Wheat
- Wild Buckwheat[5]
- Wild Mustard

Grass weeds controlled:

- Downey Brome
- Giant Foxtail
- Green Foxtail
- Persian Darnel
- Volunteer Barley
- Wild Oats

WEEDS SUPPRESSED BY PRIORITY® HL + GLYPHOSATE[1]:

- Kochia
- Annual Sow Thistle
- Perennial Sow Thistle[6]

[1] 180 g of active ingredient per acre.
[2] Less than 3 inches in height.
[3] Mature plants up to 12 inches in diameter, rosettes, and seedlings.
[4] Including all herbicide-tolerant canola varieties.
[5] Up to 5 leaves.
[6] Applications made at advanced stages will be less effective.

HOW IT WORKS:

PRIORITY® HL inhibits the production of the ALS enzyme, quickly causing plants to stop growing and become discoloured (red, yellow, purple) at the growing point and spreading to the entire plant within 1–3 weeks.

HERBICIDE

PRIORITY® HL

APPLICATION TIMING & RE-CROPPING:

- **Spring:**
  PRIORITY® HL + glyphosate may be applied in the spring prior to seeding and no longer than 48 hours after seeding prior to any crop emergence. Fields treated with PRIORITY® HL in the spring may be planted to Barley, Oats, Wheat, Durum or summer fallowed.
- **Summer:**
  Prior to August 1st: PRIORITY® HL + glyphosate may be applied to summer fallow fields and seeded in the following spring to Barley, Canola, Oats, Peas or Wheat (including durum) or summer fallowed.

  After August 1st, PRIORITY® HL + glyphosate may be applied to summer fallow fields and seeded in the following spring to Barley, Oats or Wheat (including durum) or summer fallowed.
- **Fall:**
  PRIORITY HL® + glyphosate may be applied to stubble or summer fallow fields after August 1st and prior to freeze-up and may be seeded to Barley, Oats or Wheat (including durum) or summer fallowed.

SUPPORTED TANK MIXES:

ADAMA supports the use of any glyphosate salt (DMA, IPA or K+).

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ADAMA Agricultural Solutions Canada Ltd. at 1-855-264-6262 for information before applying any tank mix that is not specifically recommended on this label.

CROP ROTATIONS:

- **When applied prior to August 1:** Barley, Canola, Oats, Field Peas and Wheat (spring, durum, winter) can be seeded the following year.
- **After August 1 and post-harvest:** Barley, Oats and Wheat (spring, durum, winter) can be seeded the following year.

PRE-HARVEST INTERVAL:

Do not harvest the treated crop within 60 days of application.

GRAZING RESTRICTIONS:

Livestock may be grazed on treated crops 7 days following application.

STORAGE:

Do not freeze, shake well before use.



CLICK HERE FOR FULL PRODUCT DETAILS.

Quick Tips

PRIORITY® HL can be mixed with the glyphosate of your choice. Remember not to mix different glyphosate salts (DMA, IPA or K+) together. Follow WAMLEGS for tank mixing directions.

ADAMA

Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

PRIORITY® HL is a registered trademark of an ADAMA Group Company.
All other products are trademarks of their respective companies.
© 2026 ADAMA Agricultural Solutions Canada Ltd.

2025-10